



Predicting Planting Stock Quality

Sylvia J. L'Hirondelle,¹ David G. Simpson,² Wolfgang D. Binder^{1,3}



B.C. Ministry of Forests and Range ¹Research Branch Laboratory, PO Box 9536 Stn Prov Govt, Victoria, BC V8W 9C4

²Kalamalka Forestry Centre, 3401 Reservoir Road, Vernon, BC V1B 2C7, ³Scientist emeritus

Storability

Stock-testing facilities need a fast, reliable method to estimate overwinter storability of seedlings (ability to survive and grow after cold storage). We compared two methods using container-grown seedlings of coastal and interior Douglas-fir, interior spruce, lodgepole pine, and western larch from British Columbia nurseries.

- Fall frost hardness was estimated using visible injury (VI) of foliage or stems and chlorophyll fluorescence (CF) of shoots after freezing to -18°C. Seedlings were then put into overwinter cold storage (-2°C).
- In spring, seedlings were planted in nursery beds; survival and growth assessed after one growing season.
- There were close correlations ($r \geq 0.93$) between VI and CF.
- Seedlings lifted after they reached 69% or higher for CF and 25% or lower for VI had over 90% survival at harvest and doubled shoot dry weight compared with seedlings lifted earlier.
- **Measuring CF was the fastest and most easily replicated method to estimate successful storability, and reduced testing time by 6 days relative to VI tests.**




Field Performance

After cold storage, conifer seedlings in British Columbia are tested for field growth potential before planting. We compared two tests of performance potential using container-grown conifer seedlings.

- Thawed seedlings were assessed for root growth potential (RGP) and chlorophyll fluorescence (CF).
- Seedlings were planted in nursery beds; survival and growth were assessed after one growing season.
- Performance tests were significantly correlated with each other ($r \geq 0.53$).
- The best performance predictor was the sum of CF + RGP ($R^2 = 0.79$ for 78 seedlot x lift-date combinations), which minimized the risk of planting poor seedlings and not planting good seedlings.
- A sum of 83 for CF + RGP provided a threshold above which survival and growth were good.
- **We recommend a combination of CF + RGP to assess vigour of shoot and root systems before planting.**

Measuring CF after freezing in the fall is useful for estimating conifer seedling storability



References:

L'Hirondelle S.J., Simpson D.G. and Binder W.D. 2006. Overwinter storability of conifer planting stock: Operational testing of fall frost hardness. *New For.* In press.

L'Hirondelle S.J., Simpson D.G. and Binder W.D. 2006. Chlorophyll fluorescence, root growth potential, and stomatal conductance as estimates of field performance potential in conifer seedlings. *New For.* In review.

Acknowledgements:

We thank current and former Ministry of Forests and Range staff for their advice and support at various stages of this project: Jim Sweeten, Clare Kooistra, Kendel Thomas, Ralph Huber, Shon Ostafew, Hazel Ritchie, B.J. Finnon, Wendy Clarke, Chris Hawkins, Glen Ursel, and Drew Brazier.

Key to abbreviations

Fdc coastal Douglas-fir	CI central interior test location
Fdi interior Douglas-fir	SI southern interior test location
Lw western larch	CS coastal test location
Pli interior lodgepole pine	CF chlorophyll fluorescence
Sx interior spruce	VI visible injury
SDW shoot dry weight	RGP root growth potential

Sum of 83 for CF + RGP in the spring seems best to minimize both risks; gives combined % survival + SDW of 140