

TASS/SYLVER/TIPSY: systems for predicting the impact of silvicultural practices on yield, lumber value, economic return and other benefits¹

(Revised September 2000)

C. Mario Di Lucca
Forest Productivity Section, Research Branch
British Columbia Ministry of Forests
Victoria, British Columbia

Abstract

Stand density management has become a very important issue in British Columbia from the point of view of timber production and its implications on economics and forest level analysis. This paper describes three systems designed to assess the affects of density management on stand yield, product recovery and quality, and financial return. Forest managers responsible for stand density decision-making can use these systems to prescribe stand-specific treatments in support of production objectives.

Introduction

This paper describes the main components and information which support three growth and yield systems used in British Columbia to predict potential yields of even-aged coniferous stands of commercial importance. The main difference between these systems is the type and complexity of information they require and produce. The Tree And Stand Simulator (TASS) generates growth and yield information for SYLVER and TIPSY. SYLVER evaluates the impact of Silviculture treatments on Yield, Lumber Value and Economic Return. It integrates the yield data from TASS and other sub-systems, and predicts wood quality, product recovery and financial return for various management regimes. Finally, TIPSY (Table Interpolation Program for Stand Yield) electronically accesses managed stand yield and product recovery tables generated by TASS and SYLVER, and performs economic analyses of the silvicultural treatments simulated.

Forest resource managers can use these systems to plan and execute stand density management prescriptions which might include different establishment densities and spatial

¹ Di Lucca, C. M. 1999. **TASS/SYLVER/TIPSY: systems for predicting the impact of silvicultural practices on yield, lumber value, economic return and other benefits.** *In:* Stand Density Management Conference: Using the Planning Tools. November 23-24, 1998, Colin R. Bamsey [Ed.] Clear Lake Ltd., Edmonton, Al. pp. 7-16.

distributions, and various silvicultural treatments (e.g. pre-commercial thinning and commercial thinning). This information can then be used for silvicultural decision-making, inventory and timber supply analysis.

The Forest Productivity Section (FPS) of the British Columbia Ministry of Forests Research Branch directs the research, development and support of these systems, with input from other ministry staff and industry. The Province of British Columbia holds the copyright for these systems.

TASS

What is TASS?

TASS is a biologically oriented spatially explicit individual tree model. It was designed to produce potential growth and yield tables for even-aged managed stands. It is calibrated for four coastal and four interior species in British Columbia. TASS is driven by the height growth (i.e. it uses the Ministry recommended site index or height-age curves), branch extension and crown expansion of competing trees. The model grows trees and simulates crown competition in a three-dimensional growing space simulated within a computer. The crowns of individual trees add a shell of foliage each year and either expand or contract asymmetrically in response to internal growth processes, physical restrictions imposed by the crowns of competitors, environmental factors and cultural practices. The volume increment produced by the foliage is distributed over the bole annually, and is accumulated to provide tree and stand statistics. More information about TASS can be found in the publications by Mitchell (1975), Mitchell and Cameron (1985) and Mitchell (1986).

TASS generates the yield table database for TIPSy. It also incorporates a number of sub-models such as ROTSIM (Mitchell and Bloomberg, 1986; Bloomberg, 1990) a laminated root-rot model, SWAT (Alfaro, 1996) a spruce weevil attack model, and TRAYCI (Brunner, 1998) a light interception model. TASS is also the growth and yield component of SYLVER which is described in the next section.

Who developed TASS?

The construction of TASS began in 1963 by Dr. Kenneth J. Mitchell as a research project of the Canadian Forestry Service (Mitchell, 1969) with later support contributed by Yale University (Mitchell, 1975), University of Idaho (Goudie, 1980), British Columbia Ministry of Forests (Mitchell and Cameron, 1985), and other agencies.

What can it do?

TASS generates growth and yield information for even-aged stands of pure coniferous species of commercial importance in coastal and interior forests of British Columbia.

It is mainly used for:

- stand level crop planning;
- stand level silvicultural treatment decision-making (e.g. espacement, fertilization, pruning, pre-commercial and commercial thinning);
- forest level planning including long-term timber supply projections for managed second-growth stands;
- pest and disease yield impact predictions (e.g. laminated root rot and spruce weevil);
- predicting height-growth repression in lodgepole pine stands;
- investigations of tree growth and stand dynamics;
- teaching growth and yield, and stand dynamics;
- generating stand density management diagrams; and
- wood quality predictions (i.e. size and distribution of branch knots and juvenile-mature wood distribution) used by SYLVER to determine product value and economic return.

The main limitation of TASS is that it does not predict the yield of complex stands (i.e. mixed-species and/or uneven-aged stands).

What data are behind TASS?

TASS is calibrated to conform with data from research trials and remeasured plots located in managed and unmanaged stands. Most of the biological growth relationships in TASS are derived from detailed stem analysis of tree boles, branches and foliage. A total of 11 989 permanent plots (i.e. comprised of 43 799 plot measurements) established for local species growing within British Columbia, Alberta, the Pacific Northwest region of the United States, Europe and New Zealand have been consistently summarized and classified by species and treatments including control, thinned and fertilized (Goudie, 1998). The number of plots and measurements from untreated natural stands and plantations is listed in Table 1.

Coastal			Interior		
Species	Number of Plots	Number of Plot Measurements	Species	Number of Plots	Number of Plot Measurements
Douglas-fir	885	4539	Lodgepole pine	2160	5351
Western hemlock	1115	4396	White spruce	456	1286
Sitka spruce	173	643	Douglas-fir	348	764
Western redcedar	38	163	Western hemlock	5	10
Total	2211	9795	Total	2969	7411

Table 1. Permanent Sample Plots Description

Ministry staff test and evaluate TASS on a regular basis using existing and new data. In a recent paper, Jim Goudie (1998) compared model output with plot data from thinned stands to identify potential biases in the predictions. He concluded that the system generally performs very well, but slightly overestimates the response to thinning.

Input options

TASS input parameters are:

- species and stand origin (i.e. single species from natural or planted origin);
- initial number of trees per hectare;
- spatial tree distribution or optional tree map list (e.g. uniform, clump or any possible distribution);
- site index or a combination of height and age;
- volume merchantability limits (e.g. minimum quadratic diameter at breast height outside bark, top diameter inside bark, and stump height);
- operational adjustment factors;
- plot dimensions in meters;
- silvicultural treatments (e.g. pre-commercial and commercial thinning, pruning and fertilization); and
- output table specifications (i.e. height or age range of the simulation and steps).

Output options

Examples of yield tables, sample output images and information produced are presented in the TASS Web site at <http://www.for.gov.bc.ca/research/gymodels/TASS>. Some of these yield tables are described below.

Standard run summary managed stand yield tables including the following variables:

- stand age;
- stand site height;
- stand crown cover percent;
- stand basal area;
- total and merchantable stand volume per hectare;
- total and merchantable stand mean annual volume increment per hectare; and
- average tree statistics (i.e. basal area, top and predominant height, crown area, crown width, crown length, quadratic mean diameter at breast height, total and merchantable height, total and merchantable volume, foliar volume, and total and merchantable gross volume).

Stand tables: display the number of trees or tree count in 5-cm diameter size classes for each stand age requested.

Stock tables: display merchantable volume (0.0+) in 5-cm diameter size classes for each stand age requested.

Mortality tables: display a stand table of the number of trees that die both earlier due to non-competitive juvenile mortality, and later due to suppression between age steps by 5-cm diameter size classes.

Standing dead tree (snag) tables: display a stand table of standing dead trees per hectare by 5-cm diameter size classes. They show the number of dead trees that are still standing at each age step (i.e. a snapshot of standing dead trees).

Customized yield tables: clients may request a variety of yield data for custom spatial distributions, pruning, fertilization, pre-commercial and commercial thinning, root rot and spruce weevil infestation treatments. Response time varies from a day to several weeks depending upon the complexity of the task.

What is in TASS's future?

Current and future developments of TASS include:

- the calibration of TRAYCI, the light interception model;
- a start-up routine to enter data from existing stands;
- a complex stands model for uneven-aged interior Douglas-fir stands in the IDF zone, and for even-aged mixed-species stands in the ICH zone;
- red alder calibration;
- internet on-line access to TASS with the support of the TASS Input Editor program (TASSIE);
- wildlife attributes and environmental indicators (e.g. pileated woodpecker, grizzly bear habitat, cougar, coarse woody debris and “old-growth” dependent values). See publication by Greenough and Kurz (1996);
- a mistletoe model for western hemlock; and
- further model calibration as data become available.

Perhaps one of the most important upgrade relates to the light and moisture components needed to simulate the development of complex stands. Light is necessary to model the variable crown structure found in mixed-species and uneven-aged stands. The dynamics of moisture within a complex stand is uncertain and is currently under investigation.

How to get TASS

TASS is only available in the Research Branch of the Ministry of Forests because it has substantial training and hardware requirements. TASS growth and yield predictions are available in two formats:

- by running the program TIPSY to obtain custom versions of the yield tables generated by TASS for a specific regime; and

- by requesting special TASS runs for yield table options that are not available within TIPSYS by contacting Ken Polsson at (250) 387-6948 or by E-mail at **Ken.Polsson@gems5.gov.bc.ca**.

In the near future, clients will submit input parameters via TASSIE over the internet.

Research Branch staff will complete the run and return the output.

More information about TASS can be found at the TASS Web site at

<http://www.for.gov.bc.ca/research/gymodels/TASS>, and in the publications by Mitchell (1969), Mitchell (1975), Mitchell (1980), Mitchell and Cameron (1985) and Mitchell (1986).

SYLVER

What is SYLVER?

SYLVER is a stand level system that helps forest managers evaluate the impact of **S**ilvicultural treatments on **Y**ield, **L**umber **V**alue, and **E**conomic **R**eturn. The components and operation of SYLVER are illustrated in Figure 1.




Figure 1: Components and Operation of SYLVER

Each element of SYLVER operates independently with information provided by the user or by the preceding elements of the system. The major components are:

- the Tree And Stand Simulator (TASS) that grows the stand and applies silvicultural treatments;
- the Bucking (BUCK) and Sawmill Simulation (SAWSIM) programs that produce logs and lumber from trees grown by TASS;
- the grading routine (GRADE) that classifies the logs and lumber according to quality; and
- the Financial Analysis System (FAN\$Y) that determines the discounted net revenue recovered from tending, harvesting and processing the stand.

Who developed SYLVER?

The development of SYLVER started in 1985 as a product of the Douglas-fir Task Force sponsored by Forintek Canada Corporation, Pulp and Paper Research Institute, British Columbia Ministry of Forests, The University of British Columbia, and industry (Kellogg, 1989). The Task Force was divided into 13 projects which studied the basic wood and pulp properties of second-growth stands of coastal Douglas-fir. The last project is the stand level system SYLVER that integrates silviculture and end-product value.

What can it do?

SYLVER predicts wood quality, product value and economic return in full for coastal Douglas-fir, and partially for coastal western hemlock, Sitka spruce, western redcedar, interior Douglas-fir, interior western hemlock, lodgepole pine and white spruce. The main applications of SYLVER are:

- stand level planning;
- silvicultural prescriptions decision-making (e.g. espacement, fertilization, pruning, pre-commercial and commercial thinning);
- forest level planning (e.g. long term timber supply projections from managed stands);
- pest and disease impact prediction (e.g. laminated root rot and spruce weevil);
- tree and stand dynamics, wood quality and economic investigations;
- stand level economics analysis; and
- education and teaching.

Some of the limitations of SYLVER are:

- it can not be distributed as an integrated stand-alone software package;
- it can not predict yield of complex stands (i.e. mixed-species and/or uneven-aged stands); and
- the lumber grade data required to calculate the select structural lumber volume is currently available only for coastal Douglas-fir.

What data are behind SYLVER?

The coastal Douglas-fir version of SYLVER is based on data from six 50-year-old stands representative of managed forests which will be harvested in the future located on Vancouver Island (Kellogg, 1989). The number of trees sampled for each project is outlined below.

- 300 trees were converted to lumber, and evaluated to estimate the log and lumber yield and grade recovery (Middleton and Munro, 1989), kiln drying degrade (Mackay, 1989), strength and stiffness of dimensional lumber (Barrett and Kellogg, 1989), and heartwood treatability (Ruddick, 1989);
- 60 trees to study the relative density distribution (Jozsa et al., 1989), chemical properties (Swan et al., 1989) and longitudinal shrinkage (Nault, 1989);
- 24 trees to study the juvenile-mature wood transition (Di Lucca, 1989);
- 17 trees to study fiber length (Hamm, 1989); and
- 9 trees to study density and chemical properties of juvenile, mature and top wood (Hatton and Hunt, 1989), unbleached Kraft pulp properties (Hatton and Hunt, 1989), and refiner mechanical pulp properties (Hatton and Johal, 1989).

Input options

The input parameters for the SYLVER components shown in Figure 1 are listed below.

- TASS input parameters were mentioned in the previous section.
- BUCK uses the tree list generated by TASS which contains the information about the tree heights, diameters inside and outside bark for the knotty and juvenile wood core, average size and distribution of knots, etc. This program bucks each tree according to specifications (i.e. log length, diameter and taper) that maximize the value of the logs or lumber.
- SAWSIM uses the log information generated by BUCK and applies a series of predefined cutting patterns to each log, and selects the one that produces the maximum value based on the determined average market prices. SAWSIM input parameters include the mill configuration (i.e. type of machine and kerf) and lumber specifications (i.e. thickness, width and length). It is a proprietary program developed by Halco Software Systems Ltd. <http://www.halcosoftware.com>.
- GRADE characterizes logs and lumber by quality class. The grading criteria for logs are: minimum length, average small-end diameter, maximum knot diameter and minimum number of annual growth rings per 2 cm. The grading criteria for clear and knotty lumber are: knot content percent, juvenile wood percent, dimensional lumber length, width and thickness.

- FAN\$Y uses stand, treatment and product information to evaluate the impact of selected silvicultural treatments on the discounted value returned by end products. Some input parameters are:
 - ◇ land manager (i.e. crown or private); land ownership (i.e. crown or private); and market type (i.e. log or lumber);
 - ◇ forestry costs (e.g. survey, land rent, brushing and weeding, spacing, fertilization, root rot control, site preparation, planting, pruning, etc.);
 - ◇ harvesting costs (e.g. infrastructure development, operation and administration overhead, tree-to-truck, hauling, etc.);
 - ◇ manufacturing costs (e.g. milling, drying, machine stress rating, etc.);
 - ◇ market prices for logs, lumber and wood chips; and
 - ◇ financial parameters (e.g. discount rate, U.S. dollar exchange rate, inflation rate, taxes, adjustments factors for future costs and prices, lumber premiums, appraisal data, etc.).

Output options

SYLVER output integrates the yield information, product recovery and financial return from a variety of management regimes. SYLVER is not distributed as a stand-alone package because TASS, BUCK and SAWSIM must be operated by specialists on a powerful workstation. In contrast, most users have only personal computers and limited training time. To provide access to SYLVER, the Forest Productivity and Decision Support Section divided the operating responsibilities between the user support service and the forest manager as illustrated in Figure 1. Both formulate a series of relevant management regimes and the specifications for bucking, sawing and grading. User support processes all regimes through to the generation of product files, which are forwarded to the forest manager for use with FAN\$Y on a personal computer. Then the forest manager can perform sensitivity analyses on the economic specifications to determine the most suitable prescription for local conditions. To contact the user support call Ken Polsson or Mario Di Lucca at (250) 387-6679 or by E-mail at **Mario.Dilucca@gems4.gov.bc.ca**.

What is in SYLVER's future?

Current and future developments of SYLVER include:

- lodgepole pine and coastal western hemlock;
- an employment generation module;
- the establishment of an expert user group for ongoing testing and development of FAN\$Y; and
- input options in TASSIE for specifying bucking, sawing and grading parameters.

How to get more information about SYLVER

More information can be found in the SYLVER Web site at <http://www.for.gov.bc.ca/research/gymodels/SYLVER>, and in the publications by Ken Mitchell, et al.(1989 and 1991). SYLVER information is also available via:

- a VHS video entitled “The SYLVER System - Forest Management Simulator” that gives a 10-minute overview of SYLVER. Copies can be borrowed by contacting Shelley Grout at (250) 387-6718 or by E-mail at Shelley.Grout@gems4.gov.bc.ca;
- a computer generated presentation called SYLVER demo (SYLDEMO) that demonstrates the importance of incorporating volume, quality and value criteria into silvicultural decisions. Currently it focuses on the operation of SYLVER to assess wood volume and value, juvenile and mature wood characteristics, and pruning by means of images, animated sequences, and supporting text. A version of this program has been modified for direct viewing at the SYLDEMO Web site at <http://www.for.gov.bc.ca/research/syldemo/syldemo>; and
- a computer program called Fred's Forest - an interactive forestry management game and training tool designed to demonstrate the capabilities of SYLVER. Players manage a woodlot for profit by selecting the regeneration method, establishment density, thinning and pruning regimes, and harvest age. It displays each treatment, and the subsequent development of the stand, with 3-dimensional graphics, followed by a score (i.e. net discounted profit). Fred's Forest is billed as a game, but its real purpose is to show the public, and kids in particular, that trees really grow and that we can manage them for our benefit. A DOS version of Fred's Forest can be downloaded from the Fred's Forest Web site at <http://www.for.gov.bc.ca/research/fredsfor>. A new multimedia CD-ROM version 3.1, which operates in both MS Windows and Macintosh environments, is now available on request from Mario Di Lucca.

TIPSY

What is TIPSY?

The **T**able **I**nterpolation **P**rogram for **S**tand **Y**ield (TIPSY) is not a growth and yield model because it only provides electronic access to the managed stand yield tables generated by TASS and SYLVER. TIPSY retrieves and interpolates yield tables from its database, customizes the information and displays summaries and graphics for a specific site, species and management regime. Information can be entered and displayed in either metric or imperial units. Initial density (i.e. planted or naturally regenerated) and pre-commercial thinning are the primary management variables. It uses optional Operational Adjustment Factors (OAFs) to mimic operational conditions. Two types of OAFs are available in TIPSY to account for elements that reduce potential yields. OAF 1 accounts for the reduction of physical growing space due to holes created by rock outcrops, swamps and non-commercial tree cover. OAF 2 accounts for pest damage that increases towards maturity. A detailed description of OAFs was prepared by Albert Nussbaum (1998). The current version of TIPSY includes an economic analysis module, known as the TIPSY Economist (Stone et al., 1996) which performs economic analyses on the silvicultural treatments simulated by TIPSY. A batch version of TIPSY is also available

for processing a large number of stands for timber supply analyses. Batch TIPSYS is included in the program WOODLOT (B.C. Ministry of Forests, 1998) for calculating even-flow harvest rates for a planning period on woodlot licenses.

Who developed TIPSYS?

The original DOS version or Meta Model was developed by Forests Planning Systems, now part of Reid Collins and Associates. In 1990, RamSoft Systems Ltd. translated the program from BASIC to the C computer language and restructured it to accommodate the next generation of yield tables. The program was renamed to TIPSYS and distributed to forest managers across the province along with an user's guide (Mitchell et al., 1992). In 1993 Ramsoft Systems Ltd. rewrote TIPSYS to operate in the MS Windows environment and renamed it WinTIPSYS. This program was released in January 1996 as Version 1.3 (Mitchell and Grout, 1995). Soon after, the development of the DOS version was discontinued and WinTIPSYS became TIPSYS again. TIPSYS Version 2.1 was released in 1998 and the current Version 3.0 was released in June 2000, and is distributed free of charge at the discretion of the Ministry.

What can it do?

TIPSYS generates managed stand yield tables, including product recovery data, economic analysis, and supporting graphics for:

- stand level crop planning;
- silvicultural treatments (e.g. espacement, fertilization, pre-commercial thinning);
- forest level planning for long term timber supply projections of managed stands. A multiple species feature aggregates up to 5 stand types into the timber supply analysis units;
- genetic gain
- commercial thinning
- repressed stands of lodgepole pine;
- dead trees (i.e. standing or fallen) and coarse woody debris;
- jobs statistics for silvicultural treatments, harvesting and manufacturing operations;
- investigations of tree growth and stand dynamic;
- generating stand density management diagrams (SDMD); and
- educational and teaching purposes.

Some of the limitations of the current version of TIPSYS are:

- it does not predict the yield of complex stands (i.e., mixed-species and/or uneven-aged stands);
- it must initiate stand growth projections at age 0 (i.e. year of disturbance). However, limited data from existing stands (i.e. top height, total number of trees, and some treatments) can be input to allow TIPSYS to estimate the number of trees at establishment;

- it allows for only one pre-commercial thinning entry, and it occurs only when stands reach a top height of 4 metres in the interior and 6 metres in the coast.
- it has only three spatial distributions. Planted stands have a nearly uniform square spacing and naturally regenerated stands have a random or clumped (i.e. for only coastal western hemlock and lodgepole pine) distribution of seedlings covering the entire site.
- it does not simulate the growth of multiple species stands biologically. The only biological assumption is the site index conversion adjustment among species.
- genetic gain is not applicable to western hemlock growing in the interior. When using the multiple species option, it's not possible to have an improved and unimproved version of the same species.
- existing stand option can be used only for stands established with 278 to 10 000 trees/ha. It can't be used for stands of hemlock (coast & interior) and lodgepole pine above 10 000 trees, even though TIPSYS database has an upper limit of 250 000 trees for these species. Consequently, it does not address repressed stands of lodgepole pine. This option is also not available for multiple species, commercial thinning or fertilization treatments.

What data are behind TIPSYS?

TIPSYS's database includes 548 managed stand yield tables for: coastal Douglas-fir, coastal western hemlock, Sitka spruce, western redcedar, interior Douglas-fir, interior western hemlock, lodgepole pine and white spruce. TIPSYS generates any table within a limited range of parameters provided by the user. If an identical yield table does not exist, TIPSYS will interpolate between the closest yield tables and electronically retrieve stand yield information.

Input options

You can enter these parameters in TIPSYS:

- species and regeneration method (e.g. natural or planted). An optional multiple species feature will prorate the yields for up to a maximum of 5 species. This option was developed to aggregate stands for the benefit of timber supply planners. It is important to understand that TIPSYS does not simulate the growth of multiple species stands biologically. The only biological assumption is the site index conversion among species;
- initial number of trees per hectare;
- operational adjustment factors;
- site index or a combination of height and age;
- silvicultural treatments (e.g. pre-commercial thinning);
- merchantability limits for total volume and merchantable volume;
- the economic analysis requires:
 - ◊ stand geography (i.e. region, district, biogeoclimatic zone and average slope);

- ◇ discount assumptions (i.e. discount rate, real price and cost increase, real increase duration, base age);
- ◇ silvicultural costs for: surveys, site preparation, planting and pre-commercial thinning;
- ◇ harvesting costs for: different harvesting systems, tree-to-truck, hauling, milling and other miscellaneous costs;
- ◇ product prices for lumber, wood chips and logs; and
- ◇ optional financial and sensitivity analyses.
- batch processing includes:
 - ◇ FIZ (Forest Inventory Zone);
 - ◇ initial density of the stand;
 - ◇ regeneration method;
 - ◇ PCT density;
 - ◇ commercial thinning;
 - ◇ operational adjustment factors 1 and 2;
 - ◇ merchantable utilization limits;
 - ◇ total area as the number of hectares that the stand occupies;
 - ◇ regeneration delay;
 - ◇ fertilization; and
- output specifications defining the beginning, end and steps of your table based on either height or age.

Output options

TIPSY generates the following yield tables and information.

Standard managed stand yield tables including:

- total age (i.e. age since disturbance);
- top height;
- gross, total and merchantable volume, and mean annual volume increment per hectare;
- stand basal area;
- quadratic mean diameter at breast height (DBH);
- crown cover percent; and
- for the largest 250 prime or crop trees: merchantable volume, quadratic mean diameter at breast height (DBH) and live crown percent.

Stand tables: display the number of trees by 5-cm diameter size classes for each stand age requested.

Stock tables: take the merchantable volume (12.5 cm+) from the yield table and display it in 5-cm diameter size classes for each stand age requested.

Stand and stock tables combined: display both the number of trees and the merchantable volume in 5-cm diameter size classes for each stand age requested.

Lumber tables: display the board-foot volume of dimensional lumber by 2-inch classes (2x4, 2x6, 2x8, and 2x10), bone dry units of chips, and lumber recovery in board feet per cubic meter.

Log tables: display the total scaled volume (12.5 cm+) of logs which are 2.5 meters (8 ft) or longer in length and its breakdown into standard log market grades.

Economic analysis tables: display the benefit/cost assumptions, net present values, site values and optional financial and sensitivity analyses of a particular management regime.

Mortality tables: display a stand table of the number of trees that die both earlier due to non-competitive juvenile mortality, and later due to suppression between age steps in 10-cm diameter size classes.

Standing dead tree (snag) tables: display a stand table of standing dead trees per hectare in 10-cm diameter size classes. They show the dead trees that are still standing at the age step (i.e. a snapshot of standing dead trees).

Coarse woody debris (CWD) tables: display the volume per hectare of CWD in the stand less the volume lost to decay, calculated by the simple exponential decay model. TIPSYS assumes that there was no CWD present on the forest floor at stand establishment. You can request the volume for All CWD created since age 0 or for CWD of a certain recruitment age in 10-cm diameter size classes.

Jobs generation tables: display the number of person days per hectare (PD/ha) of labour required for silvicultural treatments, harvesting and manufacturing operations.

Variable Density Yield Projection (VDYP) tables: display the standard VDYP output format table.

Timber Supply (TS) tables: display the standard timber supply output format table.

TIPSYS batch processing: displays three types of output tables (i.e. standard, timber supply or custom) and an error report.

TIPSYS custom tables: displays selected variables that also appear in the managed stand yield tables (yield table), log tables and lumber table. It also offers variables that do not appear in any tables, but are derived from the stand and stock table. Some examples are: average DBH, BA/ha and tree count/ha by a variety of diameter limits; and average DBH, BA/ha and merchantable volume/ha of a specified number of the trees of largest DBH (crop trees).

What is in TIPSYS's future?

The June 2000 release incorporates features completed in 1999. Future operational testing will uncover the need for a variety of improvements in addition to those already identified during very recent in-house and field testing. These improvements and new features will be added.

How to get more information about TIPSYS

More information about the TIPSYS can be found in the TIPSYS Web site at <http://www.for.gov.bc.ca/research/gymodels/TIPSYS>, and in the publication by Mitchell and Grout (1995). TIPSYS comes complete with a comprehensive on-line reference manual which includes a context-sensitive Help system as well as a directory of 'experts' who can be contacted regarding the specialized modules such as the TIPSYS Economist, snags, or lodgepole pine repression (Mitchell et al. 2000). For additional information on TIPSYS and growth and yield models, please contact Mario Di Lucca at Mario.dilucca@gems4.gov.bc.ca and for any questions regarding the installation or operation of TIPSYS, please contact Shelley Grout at shelley.grout@gems4.gov.bc.ca.

How to get TIPSYS

There are three different ways to get your copy.

- If you work for the Ministry of Forests, check with your systems administrator. All ministry offices should have recently received TIPSYS via the Data Delivery System (DDS).
- You can download TIPSYS from the Ministry of Forests Software Registration and Download Web site at <http://www.for.gov.bc.ca/research/software>.
- Finally, if you are unable to get a copy any other way, request a CD copy by contacting Shelley Grout at (250) 387-6718 or E-mail at Shelley.Grout@gems4.gov.bc.ca.

Regardless of how you get your copy of TIPSYS, you should register with the Ministry as a user. We will then be able to contact you about upgrades, bugs, and new products. If you download the software from our Web site, you will have the opportunity to register at the same time. If you do not get your copy from the Web site, you should still use our Web site to register without actually downloading.

Literature cited

Alfaro, R. I. 1992. Forecasting weevil damage. In T. Ebata (editor). Spruce Weevil Symposium Proceedings. B.C. Ministry of Forests, Prince Rupert Region, Terrace, B.C 42 pp.

- Alfaro, R. I., G. Brown, K. J. Mitchell, K. R. Polsson and R. N. Macdonald. 1996. SWAT: A decision support system for spruce weevil management. pp. 31-44. In: T L. Shore and D. L. MacLean (Eds.). Decision support systems for forest pest management. FRDA No. 260.
- Barrett, J. D. and R. M. Kellogg. 1989. Strength and Stiffness of Dimensional Lumber. In Second growth Douglas-fir: Its management and conversion for value, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 50-58.
- Bloomberg, W. J. 1990. Modeling control strategies for laminated root rot in managed Douglas-fir stands: model development. *Phytopathology*, Vol. 78, pp. 4903-4909.
- Brunner, A. In review. A light model for spatially explicit forest stand models. *For. Ecol. Manage.* 107:29-46.
- British Columbia Ministry of Forests, 1998. WOODLOT for Windows, Reference Manual. B.C. Ministry of Forests, Resource Tenure Branch. Prepared by Enfor Consultants. 18 pp.
- Di Lucca, C. M. 1989. Juvenile - Mature Wood Transition. In Second growth Douglas-fir: Its management and conversion for value, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 23-38.
- Goudie, J. W. 1980. Yield tables for managed stands of lodgepole pine in northern Idaho and southeastern British Columbia. University of Idaho, Moscow, Idaho. 111 pp.
- Goudie, J. W. 1998. Model validation: A search for the magic grove or the magic model. In Stand density management conference: Planning and implementation, November 6-7, 1997, Edmonton, Al, Bamsey, C. pp. 45-58.
- Greenough, J. A. and W. A. Kurz (1996). Stand tending impacts on environmental indicators. B.C. Min. For. Silviculture Practices Branch. FRDA II- ISBN 0-7726-2943-9. 58 pp.
- Hamm, E. 1989. Fiber Length. In Second growth Douglas-fir: Its management and conversion for value, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 44-49.
- Hatton, J. V. and K. Hunt. 1989. Density and Chemical Properties of Juvenile, Mature and Top Wood. In Second growth Douglas-fir: Its management and conversion for value, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 80-86.

Hatton, J. V. and K. Hunt. 1989. Unbleached Kraft Pulps. In *Second growth Douglas-fir: Its management and conversion for value*, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 87-113.

Hatton, J. V. and S. S. Johal. 1989. Refiner Mechanical Pulps. In *Second growth Douglas-fir: Its management and conversion for value*, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 114-129.

Jozsa, L.A., J. Richards and S. G. Johnson, 1989. Relative Density. In *Second growth Douglas-fir: Its management and conversion for value*, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 5-22.

Kellogg, R. M. (Editor), 1989. *Second growth Douglas-fir: its management and conversion for value*. Forintek Canada Corporation, Vancouver, B.C., Special Publication No. SP-32, 173 pp.

Mackay, J. F. G. 1989. Kiln Drying Lumber. In *Second growth Douglas-fir: Its management and conversion for value*, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 75-77.

Middetton, G. R. and B. D. Munro. 1989. Log and Lumber Yields. In *Second growth Douglas-fir: Its management and conversion for value*, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 66-74.

Mitchell, K. J. 1969. Simulation of the growth of even-aged stands of white spruce. In *Yale Univ. School of Forestry Bull. No 75*, Yale University, New Haven, CN. 48 pp.

Mitchell, K. J. 1975. Dynamics and simulated yield of Douglas-fir. *For. Sci. Monogr.* 17, 39 pp..

Mitchell, K. J. 1988. SYLVER: modelling the impact of Silviculture on Yield, Lumber Value and Economic Return. *For. Chron.* 64(2):127-131.

Mitchell, K. J. and I. R. Cameron. 1985. Managed stand yield tables for coastal Douglas-fir: initial density and pre-commercial thinning. B.C. Min. For., Res. Branch, Land Manage. Rep No. 31, Victoria, B.C. 69 pp.

Mitchell, K. J. and W. J. Bloomberg. 1986. Expanding concepts of growth and yield modelling to disease impacts and forest products. In *Second Can. For. Serv. Modelling Workshop, 1986*, Can. For. Serv., Victoria, B.C.

Mitchell, K. J., R. M. Kellogg and K. R. Polsson. 1989. Silvicultural Treatments and End-Product Value. In *Second growth Douglas-fir: Its management and conversion for value*. A report of the Douglas-fir Task Force, Kellogg, R. M. (eds.). Forintek Canada Corp. Spec. Publ. No. SP-32, Vancouver, B.C. pp. 130-167.

Mitchell, K. J., R. Kellogg and K. Polsson. 1989. SYLVER: The link between silvicultural practices and financial return. *Can. For. Ind.*: 38-42.

Mitchell, K. J., R. M. Kellogg and K. R. Polsson. 1991. Silvicultural treatments and end-product value. Forest Science Branch B.C. Ministry of Forests, Forest Productivity and Decision Support Section, Victoria B.C., Updated document from Mitchell et al. (1989). 38 pp.

Mitchell, K. J., S. E. Grout, R. N. Macdonald and C. A. Watmough. 1992. User's guide for TIPSYP: A Table Interpolation Program for Stand Yields, B.C. Min. For., Victoria, B.C. 78 pp.

Mitchell, K. J., and S. E. Grout. 1995. User's guide for producing managed stand yield tables for WinTIPSYP 1.3 under Microsoft Windows. B.C. Min. For., Victoria, B.C. 74 pp.

Mitchell, K. J. and J. W. Goudie. 1998. The emperor's new clothes. In *Stand density management: Planning and implementation*, November 6-7, 1997, Bamsey, C. R. (ed.). Clear Lake Ltd., Edmonton, Al. pp. 34-44.

Mitchell, K.J., M. Stone, S.E. Grout, M. Di Lucca, G.D. Nigh, J.W. Goudie, J.N. Stone, A.J. Nussbaum, A. Yanchuk, S. Stearns-Smith. R. Brockley. 2000. TIPSYP version 3.0. Online. Ministry of Forests, Research Branch, Victoria B.C. Released June 2000.

Nault, J. R. 1989. Longitudinal Shrinkage. In *Second growth Douglas-fir: Its management and conversion for value*, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 39-43.

Nussbaum, A. 1998. Operational Adjustment Factors for use in TIPSYP. Description for TFL/TSA Data Package. B.C. Min. For. Res. Br., Unp. Report, Victoria, B.C. 6 pp.

Ruddick, J. N. R. 1989. Heartwood Tretability. In *Second growth Douglas-fir: Its management and conversion for value*, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 78-79.

Stone. M. S., S. E. Grout, and C. A. Watmough. 1996. The TIPSYP Economist: An Economic Analysis module for WinTIPSYP. B.C. Min. For., Victoria. B.C. 64 pp.

Swan, E. P., J. R. Nault, C. R. Daniels and J. A. Cook. 1989. Chemical Properties. In *Second growth Douglas-fir: Its management and conversion for value*, Kellogg, R. M. (eds.). Forintek Canada Corp., Spec. Publ. No. SP-32, Vancouver, B.C. pp. 59-65.