

TABLE 5.2.1 Distribution of Fen Site Associations by biogeoclimatic zone

	BG PP	BWBS SWB	ESSF	ICH	IDF	MS	SBPS SBS	CDF	CWH	MH
Wf01 Water sedge – Beaked sedge		xx	x	xx	xxx	xxx	xxx		x ⁱ	
Wf02 Scrub birch – Water sedge		xxx	x	xx	xx	xx	xx			
Wf03 Water sedge – Peat-moss			xx				x			
Wf04 Barclay's willow – Water sedge – Glow mosses		x	xxx			x	x			
Wf05 Slender sedge – Common hook-moss		x		xx	xx	xx	xx			
Wf06 Slender sedge – Buckbean		x		x	x		x			
Wf07 Scrub birch – Buckbean – Shore sedge		x		x	x		x			
Wf08 Shore sedge – Buckbean – Hook-moss		x	x		x	x	x			
Wf09 Few-flowered spike-rush – Hook-moss			x			x	x			
Wf10 Hudson Bay clubrush – Red hook-moss							x			
Wf11 Tufted clubrush – Star moss		x	x	x		x	x			
Wf12 Narrow-leaved cotton-grass – Marsh-marigold			xxx							
Wf13 Narrow-leaved cotton-grass – Shore sedge			xx			x				
Wf50 Narrow-leaved cotton-grass – Peat-moss									x	xxx
Wf51 Sitka sedge – Peat-moss				x				xx	xx	
Wf52 Sweet gale – Sitka sedge								xx	xx ^s	
Wf53 Slender sedge – White beak-rush								x	xx ^s	

x = incidental; < 5% of wetlands

i = inland areas only

xx = minor; 5–25% of wetlands

s = southern subzones only

xxx = major; >25% of wetlands

TABLE 5.2.2 Fen Species Importance Table

Species		WF01	WF02	WF03	WF04	WF05	WF06	WF07	WF08
Shrubs	<i>Betula nana</i>								
	<i>Salix barclayi</i>								
	<i>Salix pedicellaris</i>								
	<i>Spiraea douglasii</i>								
	<i>Myrica gale</i>								
Herbs and Dwarf Shrubs	<i>Carex utriculata</i>								
	<i>Carex aquatilis</i>								
Shrubs	<i>Comarum palustre</i>								
	<i>Calamagrostis canadensis</i>								
Shrubs	<i>Carex lasiocarpa</i>								
	<i>Menyanthes trifoliata</i>								
Shrubs	<i>Carex limosa</i>								
	<i>Carex chordorrhiza</i>								
Shrubs	<i>Eleocharis quinqueflora</i>								
	<i>Trichophorum alpinum</i>								
Shrubs	<i>Trichophorum cespitosum</i>								
	<i>Eriophorum angustifolium</i>								
Shrubs	<i>Caltha leptosepala</i>								
	<i>Carex anthoxanthea</i>								
Shrubs	<i>Equisetum fluviatile</i>								
	<i>Carex magellanica</i>								
Shrubs	<i>Carex sitchensis</i>								
	<i>Rhynchospora alba</i>								
Shrubs	<i>Carex livida</i>								
	<i>Eriophorum chamissonis</i>								
Shrubs	<i>Vahlodea atropurpurea</i>								
	<i>Drosera anglica</i>								
Shrubs	<i>Hypericum anagalloides</i>								
	<i>Triantha glutinosa</i>								
Shrubs	<i>Schoenoplectus tabernaemontani</i>								
	<i>Fauria crista-galli</i>								
Shrubs	<i>Senecio triangularis</i>								
	<i>Andromeda polifolia</i>								
Shrubs	<i>Kalmia microphylla</i>								
	<i>Oxycoccus oxycoccus</i>								
Shrubs	<i>Triglochin maritima</i>								
	<i>Drosera rotundifolia</i>								
Shrubs	<i>Leptarrhena pyrolifolia</i>								
	<i>Platanthera dilatata</i>								
Shrubs	<i>Sanguisorba canadensis</i>								
	<i>Utricularia intermedia</i>								
Shrubs	<i>Viola palustris</i>								
	<i>Sphagnum</i> Group I								
Lichens and Mosses	<i>Aulaacomnium palustre</i>								
	<i>Drepanocladus</i> spp.								
Lichens and Mosses	<i>Sphagnum</i> Group II								
	<i>Tomentypnum nitens</i>								
Lichens and Mosses	<i>Philonotis fontana</i>								
	<i>Calliergon stramineum</i>								
Lichens and Mosses	<i>Scorpidium</i> spp.								
	<i>Campylopus stellatum</i>								
Lichens and Mosses	<i>Warnstorfia</i> spp.								
	<i>Meesia triquetra</i>								

Carex lasiocarpa – *Menyanthes trifoliata*

General Description

Slender sedge – Buckbean fens are uncommon in the Central and Sub-Boreal Interior at elevations below 1300 m. They occur on floating mats adjacent to small lakes and peatland ponds, or in flarks of patterned fens where there is permanent surface saturation and shallow inundation.

Sites are often slightly hummocked, with *Menyanthes trifoliata* occurring in the wet depressions and *Carex lasiocarpa* and *Drepanocladus* spp. and other mosses occurring on mounds. A sparse shrub layer can occur and the moss layer is always well developed. Hook-mosses are the most common component of the moss layer but *Sphagnum* spp. or *Campylium stellatum* may dominate on some sites.

Sites have sedge peat to 2.5 m, often with a subsurface water lens or supersaturated horizon. Fibrisols and Mesisols are typical soil types.



Characteristic Vegetation

Tree layer (0 - 0 - 0)

Shrub layer (0 - 3 - 10)

Herb layer (20 - 65 - 85)

Carex lasiocarpa, *Menyanthes trifoliata*

Moss layer (30 - 55 - 100)

Drepanocladus aduncus, *Warnstorfia* spp.

Comments

Wf06 site conditions are intermediate between the Wf05 and the Wf08. The Wf06 has a more equable water regime with less flooding, less water flow, and greater peat saturation than the related Wf05, but has deeper and more dynamic surface water than the Wf08. Slender-sedge fens (Wf05, Wf06) occur in locations similar to the Wf01 but seem to represent sites with longer surface saturation and more basic soil water. The Wf06 almost always occurs as a floating mat adjacent to a waterbody.

Similar sites in coastal areas are described by the Wf53.

Wetland Edatopic Grid




TABLE 5.3.1 Distribution of Marsh Site Associations by biogeoclimatic zone

	BG PP	BWBS SWB	ESSF	ICH	IDF	MS	SBPS SBS	CDF	CWH	MH
Wm01 Beaked sedge – Water sedge	x	xx	x	xxx	xxx	xx	xx		x	
Wm02 Swamp horsetail – Beaked sedge		x		x	x	x	xx			
Wm03 Awned sedge	x				x					
Wm04 Common spike-rush	x	x		xx	x	x	xx		x	
Wm05 Cattail	xxx	x		xx	xx	x	xx	xx	x ^s	
Wm06 Great bulrush	xxx	x		x	xx	xx	x	x	x	
Wm07 Baltic rush	x				xx					
Wm50 Sitka sedge – Hemlock-parsley								xx	xx	
Wm51 Three-way sedge				x				x	x	

x = incidental; < 5% of wetlands

xx = minor; 5–25% of wetlands

xxx = major; >25% of wetlands

s = southern subzones only

TABLE 5.3.2 Marsh Species Importance Table

Species		Wm01	Wm02	Wm03	Wm04	Wm05
Herbs and Dwarf Shrubs	<i>Carex utriculata</i>					
	<i>Carex aquatilis</i>					
	<i>Equisetum fluviatile</i>					
	<i>Comarum palustre</i>					
	<i>Sium suave</i>					
	<i>Carex exsiccata</i>					
	<i>Carex atherodes</i>					
	<i>Polygonum amphibium</i>					
	<i>Eleocharis palustris</i>					
	<i>Potamogeton richardsonii</i>					
	<i>Typha latifolia</i>					
	<i>Schoenoplectus acutus</i>					
	<i>Menyanthes trifoliata</i>					
	<i>Utricularia macrorhiza</i>					
	<i>Juncus balticus</i>					
	<i>Hordeum jubatum</i>					
	<i>Potentilla anserina</i>					
	<i>Calamagrostis canadensis</i>					
	<i>Cicuta douglasii</i>					
	<i>Lysichiton americanus</i>					
	<i>Oenanthe sarmentosa</i>					
	<i>Galium trifidum</i>					
	<i>Spiraea douglasii</i>					
	<i>Carex sitchensis</i>					
	<i>Nuphar lutea</i> ssp. <i>polysepala</i>					
	<i>Dulichium arundinaceum</i>					
Mosses	<i>Drepanocladus</i> spp.					
	<i>Wamstorfia</i> spp.					

Carex utriculata – *Carex aquatilis*

General Description

Beaked sedge – Water sedge marshes constitute the most common and widespread Marsh Site Association in the province. The **Wm01** occurs in all subzones from low to sub-alpine elevations on sites that are inundated by shallow, low-energy floodwaters and that experience some late-season drawdown. These marshes are found in a wide variety of landscape positions including flooded beaver ponds, lake margins, floodplains, and palustrine basins.



Species diversity is low and plant cover is strongly dominated by *Carex utriculata* and *C. aquatilis* with scattered forbs, aquatics, and mosses. On sites experiencing significant surface drying, species diversity increases and sites become more meadow-like. Species such as *Calamagrostis canadensis*, *Geum macrophyllum*, or *Deschampsia cespitosa* can become prominent.

The **Wm01** occurs over a wide range of site conditions on mineral substrates with thin peat veneers. Common soil types include Gleysols and Terric Humisols.

Characteristic Vegetation

- Tree layer (0 - 0 - 0)
- Shrub layer (0 - 0 - 5)
- Herb layer (13 - 80 - 100)
- Carex aquatilis*, *C. utriculata*
- Moss layer (0 - 5 - 100)

Comments

The **Wf01** and **Wm01** have similar plant communities, but, because these units are species-poor and the two dominant sedge species have a wide ecological amplitude, the plant community poorly differentiates between sites on peat (**Wf01**) and those on mineral soil (**Wm01**). In general, the **Wm01** is more deeply flooded, has more dynamic hydrology, and has a higher cover of *C. utriculata*.

The **Wm02** is another similar community that occurs on more hydrologically dynamic locations such as lake margins or floodplains. In cooler climates the **Wm01** frequently develops into **Wf01** on sites with less dynamic hydrology.

Some **Wm01** sites have scattered tall shrubs; those sites supporting > 10% shrub cover are described by Swamp Site Associations (Section 5.4).

Wetland Edatopic Grid

