

Sub Type:

Attribute Name: adjusted_ind Short Name: adjst_area
Alias adjusted ind

Forestry Term: Adjusted indicator

Description: Indicates whether or not the polygon has been adjusted.

Measurement Criteria

Standard:

Default:

Permitted Values Y - Yes; N - No

Input Format: X

Input Example: Y

Data Origin:

Attribute Source:

Sequence:
Optional:
Format: varchar2
Length: 1
Decimal Places:
Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: alpine_designation Short Name: alpn_desig
Alias: alpine designation

Forestry Term: Alpine Designation

Description: The location of the land unit with respect to location and elevation. An interpretation is applied as to whether the tree unit is above or below the tree line, that is, the upper elevation limit of continuous tree, or potential tree if cut-over, cover. If the land unit is above the the elevation line, a code of 'A' is applied, otherwise 'N', the default.

Measurement Criteria An interpretation is applied as to whether the tree unit is above or below the tree line.

Standard: If the land unit is above the elevation line, a code of 'A' is applied, otherwise 'N', the default.

Default: N

Permitted Values Codes Description

A Alpine
Alpine is the land area above the maximum elevation for tree species, dominated in vegetated areas by shrubs, herbs, bryoids and lichens. Much of the Alpine is non-vegetated covered primarily by rock, ice and snow. The Alpine is treeless by definition, however, there may be a few rare trees (<1% crown closure).

N Not Alpine
Areas not included in Alpine areas, as defined above.

Input Format: X
Input Example: N
Data Origin: input
Attribute Source: vri

Sequence:	33
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: Alpine designation contributes to the framework for delineation of ecosystems and habitat and the third level of reporting ability.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: attribution_base_date Short Name: atrib_date

Alias attribution base date

Forestry Term: Attribution Base Date

Description: The date that the information about this polygon is considered to be based on. It is currently populated, however, it is ADVISABLE to use REFERENCE DATE attribute.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: MM/DD/YYYY

Input Example:

Data Origin:

Attribute Source:

Sequence:	
Optional:	
Format:	date
Length:	
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: bark_biomass_per_ha Short Name: bioms_bark
 Alias bark_biomass_per_ha

Forestry Term: Bark Biomass

Description: this is the total whole stem biomass per hectare of all species based on a utilization of 4.0cm expressed as tonnes/ha

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
 Input Example: 235.6
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: numeric
 Length: 8
 Decimal Places: 0
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: basal_area Short Name: basal_area
 Alias basal area

Forestry Term: Basal Area at Reference Year

Description: Basal area is the total cross sectional area, at breast height, of all living trees visible to the photo interpreter in the dominant, codominant and high intermediate crown positions for each tree layer in the polygon.

Measurement Criteria The polygon is visually conceived as a whole. This impression is converted to basal area (square metres per hectare) by estimating stand structure, species composition, form factors, height by species, stems per hectare, site and uniformity.

Standard: 5 character numeric value holding basal area

Default: 0.0

Permitted Values

Input Format: #.###
 Input Example: 31.256 m2
 Data Origin: input
 Attribute Source: vri

Sequence:
 Optional:
 Format: number
 Length: 6
 Decimal Places: 3
 Null: Y

Use: Planned input for Growth Models.

Linkage:

Relationship:

Sub Type Links:

Notes: Basal area provides an estimate of polygon basal area per hectare and is used for the determination of species composition and timber volume.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: bclcs_level_1 Short Name: bclcs_lv_1
 Alias: bclcs level 1

Forestry Term: British Columbia Land Cover Classification Scheme Level 1

Description: The first level of the BC land cover classification scheme classifies the presence or absence of vegetation within the boundaries of the polygon. Presence or absence is recognized by the vertical projection of vegetation upon the land base within the polygon .

Measurement Criteria Presence or absence is recognized by the vertical projection of vegetation upon the land base within the polygon.

Standard: Level 1 is derived from the sum of the vegetation crown closures

Default:

Permitted Values

V = Vegetated
 A polygon is considered Vegetated when the total cover of trees, shrubs, herbs, and bryoids (other than crustose lichens) covers at least 5% of the total surface area of the polygon.

N = Non-Vegetated
 A polygon is considered Non-Vegetated when the total cover of trees, shrubs, herbs, and bryoids (other than crustose lichens) covers less than 5% of the total surface area of the polygon. Bodies of water are to be classified as Non-Vegetated.

U = Unreported
 A polygon is classified as Unreported if it is within the mapsheet being reported on, but is outside the inventory unit of interest. The Unreported designation is restricted to areas where inventory information is not currently available.

Examples include National Parks, Provincial Parks (where information is not available), Tree Farm Licences and Tree Farms that are not in the existing vegetation cover databases, and areas outside of the Province of British Columbia.

Note: Bodies of water may have vegetation on or under their surface; they are the responsibility of others to evaluate.

Input Format: X
Input Example: V
Data Origin: derived
Attribute Source: vri

Sequence:	36
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: The BC Land Cover Classification Scheme can be used to facilitate broad land classification reporting.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual and BC Land Classification Scheme Manual

Sub Type: veg_comp_poly

Attribute Name: bclcs_level_2 Short Name: bclcs_lv_2

Alias: bclcs level 2

Forestry Term: British Columbia Land Cover Classification Scheme Level 2

Description: The second level of the BC land cover classification scheme classifies the polygon as to the land cover type: treed or non-treed for vegetated polygons; land or water for non-vegetated polygons.

Measurement Criteria For vegetated polygons, An interpretation is made of the coverage of tree crowns as measured by their vertical projection upon the land base, estimated to the nearest percentage crown closure. For non-vegetated polygons, an interpretation is made of the percentage area occupied by land or water. For vegetated polygons an interpretation is made of the coverage of tree crowns as measured by their vertical projection upon the land base, estimated to the nearest percentage crown closure. For non-vegetated polygons, an interpretation is made of the percentage area occupied by land or water. The cover type occupying greater than 50% of the polygon area is the cover type to be assigned.

Standard: Level 2 is derived from the tree crown closure estimatem for vegetated polygons and the non-vegetated cover percent estimate for non-vegetated polygons.

Default:

Permitted Values T = Treed
A polygon is considered Treed if at least 10% of the polygon area, by crown cover, consists of tree species of any size.

N = Non-treed
A polygon is considered Non-Treed if less than 10% of the polygon area, by crown cover, consists of tree species of any size.

L = Land
The portion of the landscape not covered by water (as defined below), based on the percentage area coverage.

W = Water
A naturally occurring, static body of water, two or more metres deep in some portion, or a watercourse formed when water flows between continuous, definable banks. These flows may be intermittent or perennial; but do not include ephemeral flows where a channel with no definable banks is present. Islands within streams that have definable banks are not part of the stream; gravel bars are part of the stream. Interpretation is based on the percentage area coverage.

Input Format: X
Input Example: N
Data Origin: derived
Attribute Source: vri

Sequence:	37
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: The BC Land Cover Classification Scheme can be used to facilitate broad land classification reporting.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual and BC Land Classification Scheme Manual

Sub Type: veg_comp_poly

Attribute Name: bclcs_level_3 Short Name: bclcs_lv_3
 Alias: bclcs level 3

Forestry Term: British Columbia Land Cover Classification Scheme Level 3

Description: The location of the polygon relative to elevation and drainage, and is described as either alpine, wetland or upland. In rare cases, the polygon may be alpine wetland.

Measurement Criteria The polygon classification is determined by the landscape position with the majority coverage by area.

Standard: The Alpine designation indicates polygons that fall in the alpine regions of the landscape. For all other polygons, land cover component #1 soil moisture regime will determine whether that polygon is considered to be Upland or Wetland.

Default:

Permitted Values W = Wetland
 Land having the water table near, at, or above the soil surface, or which is saturated for a long enough period to promote wetland or aquatic processes as indicated by poorly drained soils, specialized vegetation, and various kinds of biological activity which are adapted to the wet environment.

In the Canadian wetland classification, wetland classes include bogs, fens, marshes, swamps, hot springs, hot pools, and shallow water. In British Columbia, Wetlands include forested or non-forested subhydric (SMR 7) sites, in addition to non-forested hydric (SMR 8) ecosystems (see the B.C. Land Cover Classification document for a detailed description).

U = Upland
 A broad class that includes all non-wetland ecosystems below Alpine that range from very xeric, moss- and lichen-covered rock outcrops to highly productive forest ecosystems on hygric (SMR 6) soils.

A = Alpine
 Treeless by definition (for practical purposes, 1% tree cover or less can be included within the alpine area) with vegetation dominated by shrubs, herbs, graminoids, bryoids, and lichens. Much of the Alpine is non-vegetated, covered primarily by rock, ice, and snow.

Input Format: X
Input Example: W
Data Origin: derived
Attribute Source: vri

Sequence:	38
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: The BC Land Cover Classification Scheme can be used to facilitate broad land classification reporting.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo

Interpretation Procedures Manual and BC Land Classification Scheme Manual

Sub Type: veg_comp_poly

Attribute Name: bclcs_level_4 Short Name: bclcs_lv_4
 Alias: bclcs level 4

Forestry Term: British Columbia Land Cover Classification Scheme Level 4

Description: Classifies the vegetation types and non-vegetated cover types (as described by the presence of distinct features upon the land base within the polygon).

Measurement Criteria For vegetated polygons, an interpretation is made of the coverage of vegetation crown closure as measured by their vertical projection upon the land base, estimated to the nearest percentage crown closure. For non-vegetated polygons, an interpretation is made of the non-vegetated cover percent.

Standard: If the polygon is Treed, the basal area of each species (expressed as percent composition) is reviewed and amalgamated to determine which vegetation type the polygon should be classified as. For vegetated, non-treed polygons, a hierarchical system incorporating the crown cover of shrubs, herbs and bryoids determines the Level 4 classification for the polygon.

Default:

Permitted Values TC = Treed - Coniferous
 Defined as those trees found in B.C. within the order Coniferae. These trees are commonly referred to as conifer or softwoods. The polygon is classified as Coniferous when the total basal area (expressed as percentage species composition), of coniferous trees is 75% or more of the total polygon tree basal area, and trees cover 10% or more of the total polygon area, by crown cover.

TB = Treed - Broadleaf
 Defined as those trees classified botanically as Angiospermae in the subclass Dicotyledoneae. These species are commonly referred to as deciduous or hardwoods. The polygon is classified as Broadleaf when the total basal area (expressed as percentage species composition) of broadleaf trees is 75% or more of the total polygon tree basal area, and trees cover a minimum of 10% of the total polygon area, by crown cover.

TM = Treed - Mixed
 The polygon is classified as Mixed when neither coniferous nor broadleaf trees account for 75% or more of the total polygon tree basal area, and trees cover a minimum of 10% of the total polygon area, by crown cover.

ST = Shrub Tall
 A shrub polygon with average shrub height greater than or equal to two metres.

SL = Shrub Low
 A shrub polygon with average shrub height less than two metres.

HE = Herb
 An herb polygon with no distinction between forbs and graminoids.

HF = Herb - Forbs
 An herb polygon with forbs greater than 50% of the herb cover

HG = Herb - Graminoids
 An herb polygon with graminoids greater than 50% of the herb cover.

BY = Bryoid
 A bryoid polygon with no distinction between mosses and lichens.

BM = Bryoid - Moss
 A bryoid polygon with mosses, liverworts and hornworts greater than 50% of the bryoid cover.

BL = Bryoid - Lichens
 A bryoid polygon with lichens (foliose or fruticose; not crustose) greater than 50% of the bryoid cover.

SI = Snow / Ice
 Defined as either glacier, which is considered a mass of perennial snow and ice with definite lateral limits,

typically flowing in a particular direction; or other ice and snow cover that is not part of a glacier.

RO = Rock / Rubble

Defined as bedrock or fragmented rock broken away from bedrock surfaces and moved into its present position by gravity or ice. Extensive deposits are found in and adjacent to alpine areas and are associated with steep rock walls and exposed ridges; canyons and cliff areas also contain these deposits.

EL = Exposed Land

Contains all other forms of exposed land identified by a range of subclasses.

Input Format: XX
 Input Example: TC
 Data Origin: derived
 Attribute Source: vri

Sequence:	39
Optional:	Y
Format:	varchar2
Length:	2
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: The BC Land Cover Classification Scheme can be used to facilitate broad land classification reporting.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual and BC Land Classification Scheme Manual

Sub Type: veg_comp_poly

Attribute Name:	bclcs_level_5	Short Name:	bclcs_lv_5
Alias	bclcs level 5		

Forestry Term: British Columbia Land Cover Classification Scheme Level 5

Description: Classifies the vegetation density classes and Non-Vegetated categories.

Measurement Criteria

Standard: For vegetated polygons, the Vegetation Types from Level 4 of the Scheme are further classified into density classes as listed below. Note that these are reporting breaks only and interpreters estimate density as a continuous variable. For non-vegetated polygons, the Non-Vegetated Cover Types from Level 4 of the Scheme are further classified into categories as listed below. Note that the Water cover type from Level 2 of the Scheme does not contain any classes or descriptions for water features in Level 4 of the Scheme.

Default:

Permitted Values

DE = Dense
Tree, shrub, or herb cover is between 61% and 100% for the polygon.

OP = Open
Tree, shrub, or herb cover is between 26% and 60% for the polygon.

SP = Sparse
Cover is between 10% and 25% for treed polygons, or cover is between 20% and 25% for shrub or herb polygons.

The density classes for Bryoids is as follows:

CL = Closed
Cover of bryoids is greater than 50% of the polygon.

OP = Open
Cover of bryoids is less than or equal to 50% of the polygon.

GL = Glacier
A mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction.

PN = Snow Cover
Snow or ice that is not part of a glacier but is found during summer months on the landscape.

BR = Bedrock
Unfragmented, consolidated rock, contiguous with the underlying material.

TA = Talus
Rock fragments of any size accumulated on or at the foot of slopes as a result of successive rock falls. This is a type of colluvium.

BI = Blockfield
Blocks of rock derived from the underlying bedrock by weathering and / or frost heaving. These have not undergone any significant down slope movement as they occur on level or gently sloping areas.

MZ = Rubbly Mine Spoils
Discarded overburden or waste rock, moved to extract ore during mining.

LB = Lava Bed
An area where molten rock has flowed from a volcano or fissure and cooled and solidified to form rock.

RS = River Sediments
Silt, gravel, and sand bars associated with former river channels and present river edges.

ES = Exposed Soil
Any exposed soil not covered by the other categories, such as areas of recent disturbance that include mud slides, debris torrents, avalanches, or disturbances such as pipeline rights-of-way or cultivated fields where vegetation cover is less than 5%.

LS = Pond or Lake Sediments

Exposed sediments related to dried lakes or ponds.

RM = Reservoir Margin

Land exposed by a drained or fluctuating reservoir. It is found above "normal" water levels and may consist of a range of substrates including gravel, cobbles, fine sediments, or bedrock.

BE = Beach

An area with sorted sediments reworked in recent time by wave action, which may be formed at the edge of fresh or salt water bodies.

LL = Landing

A compacted area adjacent to a road used for sorting and loading logs.

BU = Burned Area

Land showing evidence of recent burning, either natural or prescribed. Vegetation of less than 5% crown cover is present at the time of polygon description.

RZ = Road Surface

An area cleared and compacted for transporting goods and services by vehicles. Older roads that are used infrequently or not at all may cease to be classed as Non-Vegetated.

MU = Mudflat

Flat plane-like areas associated with lakes, ponds, rivers, or streams — dominated by fine-textured sediments. They can be associated with freshwater or estuarine sources.

CB = Cutbank

Part of a road corridor created upslope of the road surface, created by excavation into the hillside.

MN = Moraine

An area of debris transported and deposited by a glacier.

GP = Gravel Pit

An area exposed through the removal of sand and gravel.

TZ = Tailings

An area containing the solid waste material produced in the mining and milling of ore.

RN = Railway Surface

A roadbed with fixed rails, which may contain single or multiple rail lines.

UR = Urban

Buildings and associated developments such as roads and parking areas which form an almost continuous covering of the landscape.

AP = Airport

A permanent, paved or gravel area, and associated buildings and parking, used by airplanes.

MI = Open Pit Mine

An exposed area used to extract ore during a mining operation. This may contain associated buildings and any tailing produced by the mining and milling process.

OT = Other

A Non-Vegetated polygon where none of the above categories can be reliably chosen.

LA = Lake

A naturally occurring static body of water more than two metres deep in some portion. The boundary for the lake is the natural high water mark.

RE = Reservoir

An artificial basin affected by impoundment behind a man made structure such as a dam, berm, dyke, or wall.

RI = River/Stream

A water course formed when water flows between continuous, definable banks. Flow may be intermittent or perennial but does not include ephemeral flow where a channel with no definable banks is present. Gravel bars are part of a stream, while islands within a stream that have definable banks are not.

OC = Ocean

A naturally occurring body of water containing salt or generally considered to be salty.

Input Format:
Input Example:
Data Origin:
Attribute Source:

Sequence:	40
Optional:	Y
Format:	varchar2
Length:	2
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: The BC Land Cover Classification Scheme can be used to facilitate broad land classification reporting.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual and BC Land Classification Scheme Manual

Sub Type:

Attribute Name: branch_biomass_per_ha Short Name: bioms_brch
 Alias: branch_biomass_per_ha

Forestry Term: Branch Biomass

Description: this is the total branch biomass per hectare of all species based on a utilization of 4.0cm expressed as tonnes/ha

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
 Input Example: 235.6
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: numeric
 Length: 8
 Decimal Places: 0
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: coast_interior_cd Short Name: c_i_code
Alias: coast interior cd

Forestry Term: Coast Interior Code

Description: A code indicating that the stand is located in the Coast or Interior Region of the Province. The Coast Region is defined as the mainland west of the Cascade and Coast Mountains, including the off-shore islands. Forest Inventory Zones (FIZ) A to C are included in the Coast region. The Interior Region is defined as the mainland east of the Cascade and Coast Mountains. Forest Inventory Zones (FIZ) D to L are included in the Interior Region.

Measurement Criteria The Coast or Interior classification is used in determining stand volumes and utilization levels.

Standard: 1 character alpha code designating Coast or Interior

Default: must have value

Permitted Values I Interior (FIZ D, E, F, G, H, I, J, K and L)
C Coast (FIZ A, B, C)

Input Format: X
Input Example: C
Data Origin: derived
Attribute Source: vri

Sequence:	
Optional:	
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use: The coast or interior classifications used in determining the stand volumes and utilization levels.

Linkage:

Relationship:

Sub Type Links: vegrpt_polylayer veg_comp_poly

Notes: The Coast Region is defined as the mainland west of the Cascade and Coast Mountains, including the off-shore islands. The Interior Region is defined as the mainland east of the Cascade and Coast Mountains.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: compartment_letter Short Name: comp_let
Alias: compartment letter

Forestry Term: Inventory Compartment Letter

Description: The Compartment Letter(s) that fall within the forest cover polygon. Compartment Letter is a geographic subdivision of an Inventory Compartment. It is also part of the reference key for identifying the geographic location of all Inventory Branch samples.

Measurement Criteria: Compartment Letter only applies to some Inventory Compartments (e.g. only in Inventory Regions 1, 3, 5, 6, 7, 9, 10, 11, 56).

Standard: 1 character alpha code holding compartment letter

Default: blank

Permitted Values: <blank> No compartment Letter
A, B, C, D, E, F, G, H, I, J, K, L, M, N, P

Input Format: X
Input Example: A
Data Origin: derived
Attribute Source: both

Sequence:	
Optional:	
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use: Used to indicate the area of the polygon that is located within a compartment letter. Used in conjunction with inventory region to assign FIZ zones. Also used for defining area boundaries for area and volume summaries.

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: Compartment Letter, along with Inventory Compartment and Inventory Region form the key to identifying Inventory samples.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: crown_closure_class_cd Short Name: cc_class
Alias crown closure class code

Forestry Term: Crown Closure Class Code

Description: The class code for the ground area covered by the vertically projected crowns of the tree cover for each tree layer within the polygon.

Measurement Criteria

Standard:

Default: 0

Permitted Values
 0 0 - 5 % crown closure
 1 6 - 15 % crown closure
 2 16 - 25 % crown closure
 3 26 - 35 % crown closure
 4 36 - 45 % crown closure
 5 46 - 55 % crown closure
 6 56 - 65 % crown closure
 7 66 - 75 % crown closure
 8 76 - 85 % crown closure
 9 86 - 95 % crown closure
 10 96 - 100 % crown closure

Input Format: ##
 Input Example: 2
 Data Origin: input
 Attribute Source:

Sequence:
 Optional:
 Format: number
 Length: 2
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type: veg_comp_layer

Attribute Name: data_source_age_cd Short Name: age_dta_cd
Alias: data source age cd

Forestry Term: Data Source Age Code

Description: The source of data used for the interpretation of age and the derivation of the year of origin.

Measurement Criteria

Standard: 2 character numeric code designating method of data

Default: must have value

Permitted Values Codes Data Sources Possible Applications
0 Photo interpretation

1 Air call (air observation without 70 mm photography) species composition

2 Air call from low-level, fixed base (70 mm photography) species comp., height

3 Phase 1 photo sample (pre-1990)

4 Ground call 1 point age, height

5 Standard fixed radius sample (pre-1979) age, height

6 Phase 2 or phase 3 sample (pre-1990) species, age, height, density, basal area

7 Silviculture surveys - stocking, survival, free growing, pre-stand tending species composition, density, SMR, SNR

8 Ground observation with measurement age, height

9 Research plots (e.g. Sx trials, ecological site description) species, age, height

10 Valuation cruise plot(s) basal area, species composition, height

11 Silviculture treatment record - a record that summarizes the modified stand structure following an activity or treatment such as planting, juvenile spacing, brushing and weeding, conifer release, seed tree control, sanitation spacing, rehabilitation or commercial thinning

12 Disturbance - an area recently disturbed by fire, logging, windthrow, or insects that is classified as NSR. Has no source of information other than type and year of disturbance

13 Managed stand sample

14 Ground call, 2 or more points age, height, species composition

16 Vegetation sample age, height, density, basal area, SMR, SNR

17 Vegetation ground call age, height, density, basal area, SMR, SNR

18 Vegetation air call species composition, shrub height, shrub %

19 Natural growth sample species, age, height

20 Volume and depletion sample age, height

22 Photogrammetrically captured information that is determined or captured using photogrammetric means. An example of this is the determination of photo-measured heights using softcopy technology or parallax bars.

Input Format: ##

Input Example: 1

Data Origin:

Attribute Source:

Sequence:	52
Optional:	Y
Format:	number
Length:	2
Decimal Places:	
Null:	Y

Use: Identifies the sampling methods used to collect the layer information. The sampling method used, impacts the accuracy (e.g. confidence interval) and hence the reliability of the data

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: data_source_basal_area_cd Short Name: b_a_dta_cd
Alias: data source basal area cd

Forestry Term: Data Source Basal Area Code

Description: The source of data used for the interpretation of the basal area.

Measurement Criteria The data source will provide an indication of the reliability of attribute descriptions and may be used in the pre-inventory analysis (PIA) process. Data sources provide calibration points to aid in the determination of polygon attributes.

Standard: 2 character numeric code designating method of data

Default: must have value

Permitted Values Codes - Data Sources - Possible Applications
0 - Photo interpretation

1 - Air call (air observation without 70 mm photography) - species composition

2 - Air call from low - level, fixed base (70 mm photography) - species comp., height

3 - Phase 1 photo sample (pre - 1990) -

4 - Ground call 1 point - age, height

5 - Standard fixed radius sample (pre - 1979) - age, height

6 - Phase 2 or phase 3 sample (pre - 1990) - species, age, height, density, basal area

7 - Silviculture surveys - stocking, survival, free growing, pre - stand tending - species composition, density, SMR, SNR

8 - Ground observation with measurement - age, height

9 - Research plots (e.g. Sx trials, ecological site description) - species, age, height

10 - Valuation cruise plot(s) - basal area, species composition, height

11 - Silviculture treatment record - a record that summarizes the modified stand structure following an activity or treatment such as planting, juvenile spacing, brushing and weeding, conifer release, seed tree control, sanitation spacing, rehabilitation or commercial thinning -

12 - Disturbance - an area recently disturbed by fire, logging, windthrow, or insects that is classified as NSR. Has no source of information other than type and year of disturbance -

13 - Managed stand sample -

14 - Ground call, 2 or more points - age, height, species composition

16 - Vegetation sample - age, height, density, basal area, SMR, SNR

17 - Vegetation ground call - age, height, density, basal area, SMR, SNR

18 - Vegetation air call - species composition, shrub height, shrub %

19 - Natural growth sample - species, age, height

20 - Volume and depletion sample - age, height

22 - Photogrammetrically captured information that is determined or captured using photogrammetric means. An example of this is the determination of photo - measured heights using softcopy technology or parallax bars. - age, height

Input Format: ##
 Input Example: 3
 Data Origin: input
 Attribute Source: vri

Sequence:	32
Optional:	Y
Format:	number
Length:	2
Decimal Places:	
Null:	Y

Use: The data source may also be used to assess training issues, such as the reliability of estimates with various data sources.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: data_source_height_cd Short Name: ht_data_cd
Alias: data source height cd

Forestry Term: Data Source Height Code

Description: The source of data used for the interpretation height.

Measurement Criteria The data source will provide an indication of the reliability of attribute descriptions and may be used in the pre-inventory analysis (PIA) process. Data sources provide calibration points to aid in the determination of polygon attributes.

Standard: 2 character numeric code designating method of data

Default: must have value

Permitted Values Codes Data Sources Possible Applications
0 Photo interpretation

1 Air call (air observation without 70 mm photography) species composition

2 Air call from low-level, fixed base (70 mm photography) species comp., height

3 Phase 1 photo sample (pre-1990)

4 Ground call 1 point age, height

5 Standard fixed radius sample (pre-1979) age, height

6 Phase 2 or phase 3 sample (pre-1990) species, age, height, density, basal area

7 Silviculture surveys - stocking, survival, free growing, pre-stand tending species composition, density, SMR, SNR

8 Ground observation with measurement age, height

9 Research plots (e.g. Sx trials, ecological site description) species, age, height

10 Valuation cruise plot(s) basal area, species composition, height

11 Silviculture treatment record - a record that summarizes the modified stand structure following an activity or treatment such as planting, juvenile spacing, brushing and weeding, conifer release, seed tree control, sanitation spacing, rehabilitation or commercial thinning

12 Disturbance - an area recently disturbed by fire, logging, windthrow, or insects that is classified as NSR. Has no source of information other than type and year of disturbance

13 Managed stand sample

14 Ground call, 2 or more points age, height, species composition

16 Vegetation sample age, height, density, basal area, SMR, SNR

17 Vegetation ground call age, height, density, basal area, SMR, SNR

18 Vegetation air call species composition, shrub height, shrub %

19 Natural growth sample species, age, height

20 Volume and depletion sample age, height

22 Photogrammetrically captured information that is determined or captured using photogrammetric means. An example of this is the determination of photo-measured heights using softcopy technology or parallax bars.

Input Format: ##
Input Example: 14
Data Origin: input
Attribute Source: both

Sequence:	77
Optional:	Y
Format:	number
Length:	2
Decimal Places:	1
Null:	Y

Use: The data source may also be used to assess training issues, such as the reliability of estimates with various data sources.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes: The data source may also be used to assess training issues, such as the reliability of estimates with various data sources.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: data_src_vri_live_stem_ha_cd Short Name: stem_ha_cd
 Alias: data src vri live stem ha cd

Forestry Term: Data Source VRI Live Stem per Hectare Code

Description: The source of the data that was used for the interpretation of the vri net live stems per hectare, or stand density.

Measurement Criteria The data source will provide an indication of the reliability of attribute descriptions and may be used in the pre-inventory analysis (PIA) process. Data sources provide calibration points to aid in the determination of polygon attributes.

Standard: 2 character numeric code designating method of data

Default:

Permitted Values Codes Data Sources Possible Applications
 0 Photo interpretation

1 Air call (air observation without 70 mm photography) species composition

2 Air call from low-level, fixed base (70 mm photography) species comp., height

3 Phase 1 photo sample (pre-1990)

4 Ground call 1 point age, height

5 Standard fixed radius sample (pre-1979) age, height

6 Phase 2 or phase 3 sample (pre-1990) species, age, height, density, basal area

7 Silviculture surveys - stocking, survival, free growing, pre-stand tending species composition, density, SMR, SNR

8 Ground observation with measurement age, height

9 Research plots (e.g. Sx trials, ecological site description) species, age, height

10 Valuation cruise plot(s) basal area, species composition, height

11 Silviculture treatment record - a record that summarizes the modified stand structure following an activity or treatment such as planting, juvenile spacing, brushing and weeding, conifer release, seed tree control, sanitation spacing, rehabilitation or commercial thinning

12 Disturbance - an area recently disturbed by fire, logging, windthrow, or insects that is classified as NSR. Has no source of information other than type and year of disturbance

13 Managed stand sample

14 Ground call, 2 or more points age, height, species composition

16 Vegetation sample age, height, density, basal area, SMR, SNR

17 Vegetation ground call age, height, density, basal area, SMR, SNR

18 Vegetation air call species composition, shrub height, shrub %

19 Natural growth sample species, age, height

20 Volume and depletion sample age, height

22 Photogrammetrically captured information that is determined or captured using photogrammetric means. An example of this is the determination of photo-measured heights using softcopy technology or parallax bars.

Input Format: ##
Input Example: 3
Data Origin: input
Attribute Source: both

Sequence:	37
Optional:	Y
Format:	number
Length:	2
Decimal Places:	
Null:	Y

Use: The data source may also be used to assess training issues, such as the reliability of estimates with various data sources.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes: The data source may also be used to assess training issues, such as the reliability of estimates with various data sources.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_stand_volume_125 Short Name: dvoltot_125
 Alias: dead stand volume for 12.5 cm

Forestry Term: Dead Stand Volume for 12.5 cm

Description: This is the total net dead volume per hectare of all species determined by percent basal area of the tree layer at the 12.5 cm utilization level. Net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, net dead volumes are calculated for rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
 Input Example: 235.65
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: numeric
 Length: 8
 Decimal Places: 3
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_stand_volume_175 Short Name: dvoltot_175
 Alias: dead stand volume for 17.5 cm

Forestry Term: Dead Stand Volume for 17.5 cm

Description: This is the total net dead volume per hectare of all species determined by percent basal area of the tree layer at the 17.5 cm utilization level. Net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, net dead volumes are calculated for rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
 Input Example: 235.65
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: numeric
 Length: 8
 Decimal Places: 3
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_stand_volume_225 Short Name: dvoltot_225
 Alias: dead stand volume for 22.5 cm

Forestry Term: Dead Stand Volume for 22.5 cm

Description: This is the total net dead volume per hectare of all species determined by percent basal area of the tree layer at the 22.5 cm utilization level. Net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, net dead volumes are calculated for rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
 Input Example: 235.65
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: numeric
 Length: 8
 Decimal Places: 3
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp1_125 Short Name: dvolsp1_125

Alias dead volume per hectare for leading species at 12.5 cm

Forestry Term: Leading Species Dead Volume per Hectare at 12.5 cm

Description: This is the net dead volume per hectare of the leading species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the leading species

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp1_175 Short Name: dvolsp1_175

Alias dead volume per hectare for leading species at 17.5 cm

Forestry Term: Leading Species Dead Volume per Hectare at 17.5 cm

Description: This is the net dead volume per hectare of the leading species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the leading species

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp1_225 Short Name: dvolsp1_225

Alias dead volume per hectare for leading species at 22.5 cm

Forestry Term: Leading Species Dead Volume per Hectare at 22.5 cm

Description: This is the net dead volume per hectare of the leading species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the leading species

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp2_125 Short Name: dvolsp2_125

Alias: dead volume per hectare for second species at 12.5 cm

Forestry Term: Second Species Dead Volume per Hectare at 12.5 cm

Description: This is the net dead volume per hectare of the second species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp2_175 Short Name: dvolsp2_175

Alias: dead volume per hectare for second species at 17.5 cm

Forestry Term: Second Species Dead Volume per Hectare at 17.5 cm

Description: This is the net dead volume per hectare of the second species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp2_225 Short Name: dvolsp2_225

Alias: dead volume per hectare for second species at 22.5 cm

Forestry Term: Second Species Dead Volume per Hectare at 22.5 cm

Description: This is the net dead volume per hectare of the second species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp3_125 Short Name: dvolsp3_125

Alias: dead volume per hectare for third species at 12.5 cm

Forestry Term: Third Species Dead Volume per Hectare at 12.5 cm

Description: This is the net dead volume per hectare of the third species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp3_175 Short Name: dvolsp3_175

Alias dead volume per hectare for third species at 17.5 cm

Forestry Term: Third Species Dead Volume per Hectare at 17.5 cm

Description: This is the net dead volume per hectare of the third species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp3_225 Short Name: dvolsp3_225

Alias dead volume per hectare for third species at 22.5 cm

Forestry Term: Third Species Dead Volume per Hectare at 22.5 cm

Description: This is the net dead volume per hectare of the third species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp4_125 Short Name: dvolsp4_125

Alias: dead volume per hectare for fourth species at 12.5 cm

Forestry Term: Fourth Species Dead Volume per Hectare at 12.5 cm

Description: This is the net dead volume per hectare of the fourth species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp4_175 Short Name: dvolsp4_175

Alias: dead volume per hectare for fourth species at 17.5 cm

Forestry Term: Fourth Species Dead Volume per Hectare at 17.5 cm

Description: This is the net dead volume per hectare of the fourth species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp4_225 Short Name: dvolsp4_225

Alias: dead volume per hectare for fourth species at 22.5 cm

Forestry Term: Fourth Species Dead Volume per Hectare at 22.5 cm

Description: This is the net dead volume per hectare of the fourth species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp5_125 Short Name: dvolsp5_125

Alias: dead volume per hectare for fifth species at 12.5 cm

Forestry Term: Fifth Species Dead Volume per Hectare at 12.5 cm

Description: This is the net dead volume per hectare of the fifth species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp5_175 Short Name: dvolsp5_175

Alias: dead volume per hectare for fifth species at 17.5 cm

Forestry Term: Fifth Species Dead Volume per Hectare at 17.5 cm

Description: This is the net dead volume per hectare of the fifth species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp5_225 Short Name: dvolsp5_225

Alias: dead volume per hectare for fifth species at 22.5 cm

Forestry Term: Fifth Species Dead Volume per Hectare at 22.5 cm

Description: This is the net dead volume per hectare of the fifth species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp6_125 Short Name: dvolsp6_125
 Alias: dead volume per hectare for sixth species at 12.5 cm

Forestry Term: Sixth Species Dead Volume per Hectare at 12.5 cm

Description: This is the net dead volume per hectare of the sixth species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
 Input Example: 235.65
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: numeric
 Length: 8
 Decimal Places: 3
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp6_175 Short Name: dvolsp6_175

Alias: dead volume per hectare for sixth species at 17.5 cm

Forestry Term: Sixth Species Dead Volume per Hectare at 17.5 cm

Description: This is the net dead volume per hectare of the sixth species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:
Optional:
Format: numeric
Length: 8
Decimal Places: 3
Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: dead_vol_per_ha_spp6_225 Short Name: dvolsp6_225

Alias: dead volume per hectare for sixth species at 22.5 cm

Forestry Term: Sixth Species Dead Volume per Hectare at 22.5 cm

Description: This is the net dead volume per hectare of the sixth species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net dead volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net dead volume for the second species may be lower than volume for other species in the stand. net dead volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: earliest_nonlogging_dist_date Short Name: n_log_date
Alias: dstb_date

Forestry Term: Earliest non logging disturbance date

Description: Represents the polygons earliest non-logging disturbance date.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: MM/DD/YYYY

Input Example:

Data Origin:

Attribute Source:

Sequence:
Optional:
Format:
Length:
Decimal Places:
Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: earliest_nonlogging_dist_type Short Name: n_log_dist
 Alias: dstb_type

Forestry Term: Earliest non logging disturbance type

Description: Represents the polygons earliest non-logging disturbance type.

Measurement Criteria

Standard: 3 character code

Default:

Permitted Values This is a 3 character field.
 The first character represents the following:
 •General Disturbance
 The second and third character describes the non-logging disturbance type.

- General Disturbance
 ++++++
- A - Animal damage (general) DI
 - B - Wildfire DI
 - BE - Escaped burn (DI)
 - BG - Ground fire (DI)
 - BR - Range burn (DI)
 - BW - Wildlife burn (DI)
 - D - Diseases (general) (DI)
 - F - Flooding (DI)
 - I - Insects (general) (DI)
 - K - Fume Kill (DI)
 - L - Logging (DI)
 - N - Non-Biological (abiotic) injuries (DI)
 - R - Site rehabilitation (DI)
 - S - Slide or Avalanche (DI)
 - T - Treatment injuries (general) (DI)
 - U - Damage (cause unknown) (DI)
 - V - Problem vegetation (DI)
 - W - Windthrow (DI)

+++++

- A -Animal Damage (general)
 ++++++
- B - Bear
 - C - Cattle
 - D - Deer
 - E - Elk
 - H - Hare or Rabbit
 - M - Moose
 - P - Porcupine
 - S - Squirrel
 - V - Vole
 - X - Birds
 - Z - Beaver

- D - Diseases (general)
 ++++++
- A - Foliages Diseases (general)
 - AF - Broom Rusts
 - B - Brooming (Non-Mistletoe)
 - BF - Fir Broom Rust
 - BS - Spruce Broom Rust
 - D - Stem Rots (general)
 - DA - Armillaria
 - DC - Laminated, Cedar Strain Only

DE - Rust-Red Stringy Rot
 DF - Brown Crumbly Rot
 DI - Tomentosus
 DL - Laminated (not cedar strain)
 DN - Annosus
 DP - Red Ring Rot
 DS - Schweinitzii Butt Rot
 DT - Aspen Trunk Rot
 F - Foilage Diseases (general)
 FB - Larce Needle Blight
 FD - Douglas-fir Needle Cast
 FE - Elytroderma Needle Cast
 FL - Lophodermelia (Pine) Needle Cast
 FM - Larch Needle Cast
 FP - Fir Needle Blight (Fir-fireweed Rust)
 FR - Red Band Needle (Blight) Cast
 L - Leader and Branch Dieback (general)
 LD - Derma Canker
 LL - Leader Dieback
 LP - Phomopsis Canker
 LR - Branch Dieback
 LS - Sydowia (Sclerophoma) Tip Dieback
 LV - Aspen-poplar Twig Blight
 M - Dwarf Mistletoe (general)
 MF - Douglas-fir Dwarf Mistletoe
 MH - Hemlock Dwarf Mistletoe
 ML - Larch Dwarf Mistletoe
 MP - Lodgepole Pine Dwarf Mistletoe
 P - Bark Disease (general)
 R - Root Disease (general)
 RA - Amarillaria Root Disease
 RB - Black Stain Root Disease
 RC - Laminated Root Rot, Cedar Strain
 RL - Laminated Root Rot, not Cedar Strain
 RN - Annosus Root Rot
 RR - Rhizina Root Disease
 RT - Tomentosus Root Rot
 S - Stem Diseases (general)
 SA - Atropellis Canker (Lodgepole Pine)
 SB - White Pine Blister Rust
 SC - Comandra Blister Rust
 SG - Western Gall Rust
 SN - Aspen Cankers (Cystopora Canker)
 SN - Aspen Cankers (Hypoxylon Canker)
 SN - Aspen Cankers (Cryptospheria Canker)
 SN - Aspen Cankers (Target Canker)
 SN - Aspen Cankers (Ceratocystis Canker)
 SN - Aspen Cankers (Sooty Bark Canker Canker)
 SS - Stalactiform Blister Rust
 SX - Exploding Canker of Douglas-fir and Interior Spruces

I - Insects (general)
 ++++++

A - Aphids (general)
 AB - Balsam Woolly Adelgid
 AC - Gian Conifer Aphid
 AG - Colley Spruce Gall Adelgid
 AS - Green Spruce Aphid
 B - Bark Beetles (general)
 BB - Western Balsam Bark Beetle
 BD - Douglas-fir Beetle
 BI - Engraver Beetle
 BM - Mountain Pine Beetle
 BP - Twig Beetle and Others
 BS - Spruce Beetle
 BT - Red Turpentine Beetle
 BW - Western Pine Beetle
 D - Defoliations (general)
 DA - Black Army Cutworm
 DC - Larch Casebearer

- DD - Loper (Deciduous)
- DE - Eastern Spruce Budworm
- DF - Forest Tent Caterpillar
- DG - Greenstriped Forest Looper
- DH - Western Blackheaded Budworm
- DM - Gypsy Moth
- DN - Birch Leaf Miner
- DP - Larch Sawfly
- DR - Red Alder Sawfly
- DS - Conifer Sawfly
- DT - Douglas-fir Tussock Moth
- DU - Satin Moth
- DV - Variegated Cutworm
- DW - Western Aspen Tortrix
- DZ - Western False Hemlock Looper
- M - Mite Damage (general)
- S - Shoot Insects (general)
- SB - Western Cedar Borer
- SE - European Pine Shoot Moth
- SG - Gouty Pitch Midge
- SP - Pitch Nodule Moths
- SQ - Sequoia Pitch Moth
- SS - Western Pine Shoot Borer
- W - Weevils (general)
- WC - Steremnius Root Collar Weevil
- WM - Magdalis Species
- WP - Lodgepole Pine Terminal Weevil
- WS - White Pine (Spruce) Weevil
- WW - Warren's Root Collar Weevil
- WY - Cyliindrocopturus Weevil
- WZ - Yosemite Bark Weevil

- N - Non-Biological (abiotic) Injuries
 ++++++
- B - Wildfire
- D - Drought
- F - Flooding
- G - Frost (general)
- GC - Frost Crack
- GH - Frost Heaved
- GK - Shoot/Bud Frost Kill
- Hv - Hail
- K - Fume Kill
- L - Lightning
- N - Road Salt
- R - Redbelt
- S - Slide
- W - Windthrow
- WS - Windthrow-soil failure
- WT - Windthrow-treatment or harvest related
- X - Scarring/rubbing
- Y - Snow or Ice (including Snow Press)
- Z - Sunscald

- T - Treatment Injuries (general)
 ++++++
- C - Chemical
- L - Logging
- M - Mechanical
- P - Planting
- PM - Poor Planting Microsite
- R - Pruning
- T - Thinning or Spacing

- U - Unknown Damage (cause Unknown)
 ++++++
- A - Atypical Growth
- B - Breakage (Dead or Broken Top)
- C - Crown Symptoms (Chlorotic)
- F - Fluted Butt

G - Galls
 K - Fork or Pronounced Crook
 L - Leaning
 M - Multiple Leaders
 R - Brooming
 S - Basal Sweep

V - Problem Vegetation
 ++++++

H - Herbaceous Competition
 P - Vegetation Press
 S - Shrub Competition
 T - Tree Competition

Input Format: XXX

Input Example:

Data Origin:

Attribute Source:

Sequence:
Optional:
Format: varchar2
Length: 2
Decimal Places:
Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name:	ecosys_class_data_src_cd	Short Name: eco_src_cd
Alias	ecosys class data src cd	
Forestry Term:	Ecosystem Class Data Source Code	
Description:	The source of the data used in the interpretation of the ecological attributes (Surface expression, modifying process, site position meso, alpine designation, and soil nutrient regime) that describe the polygon.	
Measurement Criteria	The data source will provide an indication of the reliability of attribute descriptions and may be used in the pre-inventory analysis (PIA) process. Data sources provide calibration points to aid in the determination of polygon attributes.	
Standard:	2 character numeric code designating method of data	
Default:		
Permitted Values	<p>Codes Data Sources Possible Applications</p> <p>0 Photo interpretation</p> <p>1 Air call (air observation without 70 mm photography) species composition</p> <p>2 Air call from low-level, fixed base (70 mm photography) species comp., height</p> <p>3 Phase 1 photo sample (pre-1990)</p> <p>4 Ground call 1 point age, height</p> <p>5 Standard fixed radius sample (pre-1979) age, height</p> <p>6 Phase 2 or phase 3 sample (pre-1990) species, age, height, density, basal area</p> <p>7 Silviculture surveys - stocking, survival, free growing, pre-stand tending species composition, density, SMR, SNR</p> <p>8 Ground observation with measurement age, height</p> <p>9 Research plots (e.g. Sx trials, ecological site description) species, age, height</p> <p>10 Valuation cruise plot(s) basal area, species composition, height</p> <p>11 Silviculture treatment record - a record that summarizes the modified stand structure following an activity or treatment such as planting, juvenile spacing, brushing and weeding, conifer release, seed tree control, sanitation spacing, rehabilitation or commercial thinning</p> <p>12 Disturbance - an area recently disturbed by fire, logging, windthrow, or insects that is classified as NSR. Has no source of information other than type and year of disturbance</p> <p>13 Managed stand sample</p> <p>14 Ground call, 2 or more points age, height, species composition</p> <p>16 Vegetation sample age, height, density, basal area, SMR, SNR</p> <p>17 Vegetation ground call age, height, density, basal area, SMR, SNR</p> <p>18 Vegetation air call species composition, shrub height, shrub %</p> <p>19 Natural growth sample species, age, height</p> <p>20 Volume and depletion sample age, height</p> <p>22 Photogrammetrically captured information that is determined or captured using photogrammetric means. An example of this is the determination of photo-measured heights using softcopy technology or parallax bars.</p>	

Input Format: ##
Input Example: 3
Data Origin: input
Attribute Source:

Sequence:	35
Optional:	Y
Format:	number
Length:	2
Decimal Places:	
Null:	Y

Use: The data source may also be used to assess training issues, such as the reliability of estimates with various data sources.

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: est_coverage_pct_2 Short Name: cov_pct_2

Alias land cover component percentage 2

Forestry Term: Land Cover Component Percentage 2

Description: The amount the polygon occupied by the second most dominate Land Cover Component. The sub-division of a polygon by a quantified Land Cover Component allows a higher degree spatial resolution for modeling wildlife habitat capability. Generally, sizes under 10% would not be estimated.

Measurement Criteria Examine the polygon to determine how many land cover components are present. Individual component pieces that make up less than 10% of the polygon area should not be estimated for land cover components. Each land cover component should have a minimum of one contiguous piece that makes up at least 10% of the polygon area. Individual pieces or patches that are individually at least 10% of the polygon may be combined such that the amalgamation of those individual pieces could constitute 20% of the polygon area or greater. Record land cover component percent to the nearest percent. The total of all land cover component percent values must equal 100%.

Standard: 3 charater numeric

Default:

Permitted Values 0 to 100

Input Format: ###

Input Example: 10

Data Origin: input

Attribute Source:

Sequence:
 Optional:
 Format: number
 Length: 3
 Decimal Places:
 Null:

Use: Land cover component percent quantifies the extent of each land cover component identified. It provides for reporting to a finer resolution than the polygon unit and can be used to model wildlife habitat capability.

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: est_coverage_pct_3 Short Name: cov_pct_3

Alias land cover component percentage 3

Forestry Term: Land Cover Component Percentage 3

Description: The amount the polygon occupied by the third most dominate Land Cover Component. The sub-division of a polygon by a quantified Land Cover Component allows a higher degree spatial resolution for modelling wildlife habitat capability. Generally, sizes under 10% would not be estimated.

Measurement Criteria Examine the polygon to determine how many land cover components are present. Individual component pieces that make up less than 10% of the polygon area should not be estimated for land cover components. Each land cover component should have a minimum of one contiguous piece that makes up at least 10% of the polygon area. Individual pieces or patches that are individually at least 10% of the polygon may be combined such that the amalgamation of those individual pieces could constitute 20% of the polygon area or greater. Record land cover component percent to the nearest percent. The total of all land cover component percent values must equal 100%.

Standard: 3 charater numeric

Default:

Permitted Values 0 to 100

Input Format: ###

Input Example: 10

Data Origin: input

Attribute Source:

Sequence:
 Optional:
 Format: number
 Length: 3
 Decimal Places:
 Null:

Use: Land cover component percent quantifies the extent of each land cover component identified. It provides for reporting to a finer resolution than the polygon unit and can be used to model wildlife habitat capability.

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: est_coverage_pct_1 Short Name: cov_pct_1
 Alias land cover component percentage 1

Forestry Term: Land Cover Component Percentage 1

Description: The amount the polygon occupied by the predominate Land Cover Component. The sub-division of a polygon by a quantified Land Cover Component allows a higher degree spatial resolution for modelling wildlife habitat capability. Generally, sizes under 10% would not be estimated.

Measurement Criteria Examine the polygon to determine how many land cover components are present. Individual component pieces that make up less than 10% of the polygon area should not be estimated for land cover components. Each land cover component should have a minimum of one contiguous piece that makes up at least 10% of the polygon area. Individual pieces or patches that are individually at least 10% of the polygon may be combined such that the amalgamation of those individual pieces could constitute 20% of the polygon area or greater. Record land cover component percent to the nearest percent. The total of all land cover component percent values must equal 100%.

Standard: 3 character numeric

Default:

Permitted Values 0 to 100

Input Format: ###
 Input Example: 10
 Data Origin: input
 Attribute Source:

Sequence:
 Optional:
 Format: number
 Length: 3
 Decimal Places:
 Null:

Use: Land cover component percent quantifies the extent of each land cover component identified. It provides for reporting to a finer resolution than the polygon unit and can be used to model wildlife habitat capability.

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: est_site_index Short Name: est_si
Alias: est site index

Forestry Term: Estimated Site Index

Description: Estimated site index is an interpreter estimated site index for tree layers with a leading species age less than 31 years. Site index is the mean height of the dominant and codominant trees will attain at a base index age (50 years) used for the purposes of estimating forest site growth capability. The site index is based on a normalized set of coefficients calibrated to reflect the range of heights for a given tree species.

Measurement Criteria Estimated site index may be based on the direct application of conventional site index curves, or it may be estimated from other data sources. The direct site index value may be determined from the dominant and codominant trees.

Standard: 2 character numeric value holding estimated site index in metres (bha 50). Site index estimates are required on all treed polygons as well as polygons that are potentially capable of producing trees.
2 character numeric value holding estimated site index in metres (bha 50).

Default:

Permitted Values

Input Format: ##
Input Example: 15
Data Origin: input
Attribute Source: both

Sequence:	14
Optional:	Y
Format:	number
Length:	2
Decimal Places:	
Null:	Y

Use: Used in assigning culmination MAI for young stands in Timber Supply Analyses and Local Resource Use Plans (LRUPs). Also used as a basis for applying net-downs for low sites.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes: Used in assigning Culmination MAI for young stands in Timber Supply Analyses and Local Resource Use Plans (LRUPs). Also used as a basis for applying net-downs for low sites.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: est_site_index_source_cd Short Name: si_data_cd
 Alias estimates site index source code

Forestry Term: Estimates site index source code

Description: Describes the process used to determine the estimated site index prediction. for tree layers with a leading species age less than 30 years

Measurement Criteria Estimates site index source indicates the method used for obtaining an estimated site index, the reliability of the estimate and classifies the sources for the analysis.

Standard:

Default:

Permitted Values
 A Adjacent stand
 The site index is assigned using information from adjacent stands with similar species, age and height.
 C Site Index Curve
 E Ecological correlation
 Using an assessment of ecological site factors and indicator plant species prior to harvest, a determination is made of an ecological classification. The classification and associated site index for various species is attained from tabular values.
 H Historic
 Derived from the site index value of the previous stand with no change to the site index value.
 I Growth intercept
 This is a field procedure carried out on stands that have at least five years growth above breast height, but are less than 30 years old. It is determined during a silvicultural survey.
 M Site Class Conversion
 O SIBEC rollover
 S Silviculture section
 Assigned by the District Silviculture section; when the method of determination is unknown (potential methods can be the growth intercept, ecological correlations, historic, or adjacent stands).

Input Format: X
 Input Example: A
 Data Origin: input
 Attribute Source:

Sequence:
 Optional:
 Format: varchar2
 Length: 1
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: est_site_index_species_cd Short Name: est_si_spc
Alias: est site index species cd

Forestry Term: Estimated Site Index Species Code

Description: Estimated site index species is the tree species from which the site index for the polygon has been estimated. The site index species provides a link between the estimated site index and a particular tree species' productivity at that site.

Measurement Criteria: The interpreter will view the polygon and select the tree species that provides the best description of site productivity.

Standard: For polygons with trees less than 30 years (total age); currently non-treed but capable of producing trees; and occupied by trees planted outside their normal ecological range.

Default:

Permitted Values

NATIVE CONIFERS ---
Cedar -Thuja -C -
western redcedar -Thuja plicata - -Cw

Cypress -Chamaecyparis -Y -
yellow-cedar -C. nootkatensis - -Yc

Douglas-fir -Pseudotsuga -F -
Douglas-fir -P. menziesii - -Fd
coastal Douglas-fir -P. menziesii var. menziesii - -Fdc
interior Douglas-fir -P. menziesii var. glauca - -Fdi

Fir (Balsam) -Abies -B -
amabilis fir -A. amabilis - -Ba
grand fir -A. grandis - -Bg
subalpine fir -A. lasiocarpa - -Bl

Hemlock -Tsuga -H -
mountain hemlock -T. mertensiana - -Hm
western hemlock -T. heterophylla - -Hw
mountain x western hemlock hybrid -T. mertensiana x heterophylla - -Hxm

Juniper -Juniperus -J -
Rocky Mtn. juniper -J. scopulorum - -Jr

Larch -Larix -L -
alpine larch -L. lyallii - -La
tamarack -L. laricina - -Lt
western larch -L. occidentalis - -Lw

Pine -Pinus -P -
jack pine -P. banksiana - -Pj
limber pine -P. flexilis - -Pf
lodgepole pine -P. contorta - -Pl
lodgepole pine -P. contorta var. latifolia - -Pli
lodgepole x jack pine hybrid -P. x murraybanksiana - -Pxi
ponderosa pine -P. ponderosa - -Py
shore pine -P. contorta var. contorta - -Plc
western white pine -P. monticola - -Pw
whitebark pine -P. albicaulis - -Pa

NATIVE CONIFERS ---
Spruce -Picea -S -
black spruce -P. mariana - -Sb
Engelmann spruce -P. engelmannii - -Se
Sitka spruce -P. sitchensis - -Ss
white spruce -P. glauca - -Sw
spruce hybrid -Picea cross - -Sx
Engelmann x white -P. engelmannii x glauca - -Sxw

Sitka x white -P. x lutzii - -Sxl
Sitka x unknown hybrid -P. sitchensis x ? - -Sxs

Yew -Taxus -T -
western yew -Taxus brevifolia - -Tw

NATIVE HARDWOODS ---

Alder -Alnus -D -
red alder -A. rubra - -Dr

Apple -Malus -U -
Pacific crab apple -Malus fusca - -Up

Aspen, Cottonwood or Poplar -Populus -A -
poplar -P. balsamifera - -Ac
balsam poplar -P. b. ssp. balsamifera - -Acb
black cottonwood -P. b. ssp. trichocarpa - -Act
hybrid poplars -P. spp. - -Ax
trembling aspen -P. tremuloides - -At

Arbutus -Arbutus -R -
Arbutus -Arbutus menziesii - -Ra

Birch -Betula -E -
Alaska paper birch -B. neoalaskana - -Ea
Alaska x paper birch hybrid -B. x winteri - -Exp
paper birch -B. papyrifera - -Ep
water birch -B. occidentalis - -Ew

Cascara -Rhamnus -K -
cascara -R. purshiana - -Kc

Cherry -Prunus -V -
bitter cherry -P. emarginata - -Vb
choke cherry -P. virginiana - -Vv
pin cherry -P. pensylvanica - -Vp

Dogwood -Cornus -G -
Pacific dogwood -Cornus nuttallii - -Gp

Maple -Acer -M -
bigleaf maple -A. macrophyllum - -Mb
vine maple -A. circinatum - -Mv

Oak -Quercus -Q -
Garry oak -Q. garryana - -Qg

Willow -Salix spp. -W -
Bebb's willow -S. bebbiana - -Wb
Pacific willow -S. lucida - -Wp
peachleaf willow -S. amygdaloides - -Wa
pussy willow -S. discolor - -Wd
Scouler's willow -S. scouleriana - -Ws
Sitka willow -S. sitchensis - -Wt

UNKNOWNNS ---

Unknown - -X -
Unknown conifer - - -Xc
Unknown hardwood - - -Xh

OTHERS ---

Other tree, not on list - -Z -
Other conifer - - -Zc
Other hardwood - - -Zh

EXOTICS ---

Apple -Malus -U -
apple -Malus pumila - -Ua
Aspen, Cottonwood or Poplar -Populus -A -
*southern cottonwood -P. deltoides - -Ad

Birch -Betula -E -
 European birch -B. pendula - -Ee
 silver birch -B. pubescens - -Es
 Cherry -Prunus -V -
 sweet cherry -P. avium - -Vs
 Cypress -Chamaecyparis -Y -
 *Port Orford-cedar -C. lawsoniana - -Yp

EXOTICS ---
 Fir (Balsam) -Abies -B -
 *balsam fir -A. balsamea - -Bb
 noble fir -A. procera - -Bp
 *Shasta red fir -A. magnifica var. shastensis - -Bm
 *white fir -A. concolor - -Bc

Maple -Acer -M -
 box elder -A. negundo - -Me
 *Norway maple -A. platanoides - -Mn
 *Sycamore maple -A. pseudoplatanus - -Ms

Other exotics - - -
 *incense-cedar -Calocedrus decurrens - -Oa
 *giant sequoia -Sequoiadendron giganteum - -Ob
 *coast redwood -Sequoia sempervirens - -Oc
 European mountain-ash -Sorbus aucuparia - -Od
 Siberian elm -Ulmus pumila - -Oe
 common pear -Pyrus communis - -Of
 Oregon ash -Fraxinus latifolia - -Og

Pine -Pinus -P -
 *Monterey pine -P. radiata - -Pm
 *red pine -P. resinosa - -Pr
 *sugar pine -P. lambertiana - -Ps
 Oak -Quercus -Q -
 *English oak -Q. robur - -Qe
 Spruce -Picea -S -
 *Norway spruce -P. abies - -Sn

Input Format: XX
 Input Example: HX
 Data Origin: input
 Attribute Source: both

Sequence:	13
Optional:	Y
Format:	varchar2
Length:	3
Decimal Places:	
Null:	Y

Use:
 Linkage:
 Relationship:
 Sub Type Links: veg_comp_layer

Notes: The estimate of site index species provides a link between the site index and a particular tree species site productivity (i.e., age / height curve).

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo

Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: feature_class_skey Short Name: feat_skey
Alias: feature class skey

Forestry Term: Feature Class Skey

Description: Unique identifier for a feature class.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin: generated

Attribute Source:

Sequence:	
Optional:	
Format:	number
Length:	38
Decimal Places:	
Null:	N

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: feature_id Short Name: feature_id
Alias feature identity

Forestry Term: Feature Identity

Description: Provincially unique identifier for an instance of a spatial feature

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	
Optional:	
Format:	varchar2
Length:	32
Decimal Places:	
Null:	N

Use:

Linkage:

Relationship:

Sub Type Links: vegrpt_polylayer veg_comp_layer veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: fiz_cd Short Name: fiz_cd
 Alias fiz cd

Forestry Term: Forest Inventory Zone

Description: The Forest Inventory Zone(s) (FIZ) that fall within the forest cover polygon. FIZ zones were developed to provide a broadly based ecological classification of the forestland in British Columbia. FIZ zones closely follow the early Biogeoclimatic zones developed by Dr. Krajina. The province of British Columbia is split into 12 FIZ zones.

Measurement Criteria

Standard: 1 character alpha code holding FIZ (A to L)

Default: must have value

Permitted Values A to L

Input Format: X
 Input Example: K
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: varchar2
 Length: 1
 Decimal Places:
 Null: Y

Use: Used to indicate the area of the polygon located within a FIZ zone. Used in conjunction with Public Sustained Yield Unit to assign stand volumes.

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: Used to indicate the area of the polygon located within a FIZ zone. Used in conjunction with Public Sustained Yield Unit to assign stand volumes.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: foliage_biomass_per_ha Short Name: bioms_fol
Alias: foliage_biomass_per_ha

Forestry Term: Folliage Biomass

Description: this is the total bark biomass per hectare of all species based on a utilization of 4.0cm expressed as tonnes/ha

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.6

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 0

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: for_cover_rank_cd Short Name: rank_cd
 Alias for cover rank cd

Forestry Term: Forest Cover Rank Code - Rank or Importance of Layer

Description: A numeric designation of the relative importance of the layer component in the stand as determined by the business. For VRI originated data, this value is assigned via business rule based on the supplied order of the layer records as recorded by the interpreter. For FIP originated data, this value is known as the RANK CD, and is explicitly supplied by the interpreter. The RANK CD, or ranking, was based on Regional guidelines at the time of interpretation. This value is retained for FIP transition purposes, as tree volumes are only calculated by VDYP, the current software/mathematical model in production. The RANK CD determines which layer will provide attributes to the yield prediction model (VDYP).

Measurement Criteria One layer in a multi-layered stand is assigned a rank code indicating the relative importance of that layer. The layer assigned Rank 1 in multi-layer stands is the most important layer

Standard: For Vegetation Cover originated data, this value is assigned via a series of business rules based on the species composition, age, height and crown closure of the layers as recorded by the interpreter.

Default: blank

Permitted Values 1 Rank 1, most important layer

Input Format: #
 Input Example: 2
 Data Origin: input / derived
 Attribute Source: fip / vri

Sequence:	8
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use: Defines the importance of the layer. Only Rank 1 layers are used when summarizing the land base for Timber Supply Analyses. RANK 1 is used to define the label description for Forest Cover Maps.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes: Volumes are calculated for Rank 1 stands only. Rank assignment is based on Regional guidelines.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: for_mgmt_land_base_ind Short Name: fmlb
Alias forest management land base

Forestry Term: Forest Management Land Base

Description: An indicator placing the polygon in the Forest management Land Base. This is the equivalent to THLB (as a starting point)

The criteria is if site index >5, opening_ind (opening) equal to "Y" (yes), or inventory_standard_cd equal V or I but bclcs_level_1 not equal "N", or bclcs_level_3 not equal "A"

Measurement Criteria

Standard:

Default:

Permitted Values Y - Yes; N - No

Input Format: X
Input Example: Y
Data Origin: derived
Attribute Source: both

Sequence:
Optional:
Format: varchar2
Length: 1
Decimal Places:
Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: free_to_grow_ind Short Name: ftg_ind
Alias free_to_grow_ind

Forestry Term: Free to grow indicator

Description: Indicates whether or not the polygon represents a Free To Grow opening, a point where the basic silviculture obligations have been fulfilled.

Measurement Criteria

Standard:

Default:

Permitted Values Y - Yes; N - No

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:

Optional:

Format: varchar2

Length: 1

Decimal Places:

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: full_label Short Name: full_label
 Alias full label

Forestry Term: Full Label

Description: The full Vegetation Map label. It contains the polygon id, opening number, species composition, projected age class, projected height class, crown closure class, a hyphen, site index, a forward slash, estimated site index, and code(s) for shrub, herb, bryoid, or non vegetative components, and the historic disturbance and forest management activities. It is at most 8 lines. Back slashes represent carriage returns.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	
Optional:	
Format:	varchar2
Length:	500
Decimal Places:	
Null:	N

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: geometry Short Name: geometry
Alias geometry

Forestry Term: Geometry

Description: A spatial polygon feature used to store the map component of the Vegetation Cover area.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source: calculated and generated

Sequence:
Optional:
Format: number
Length:
Decimal Places:
Null: N

Use:

Linkage:

Relationship:

Sub Type Links: veg_vegetation_cover_polygon

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: harvest_date Short Name: harv_date

Alias harv_date

Forestry Term: Harvest Date

Description: The date in which the polygon was last harvested.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: MM/DD/YYYY

Input Example:

Data Origin:

Attribute Source: Results

Sequence:
Optional:
Format: date
Length:
Decimal Places:
Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: herb_cover_pattern Short Name: herb_cover
 Alias herb cover pattern

Forestry Term: Herb Cover Pattern

Description: Herb cover pattern is a code that describes the spatial distribution of the herbaceous species within the polygon. Herb cover pattern is used to describe the herb layer spatial distribution. Examples include clumps of herbaceous species on rock outcrops, scattered patches or individual herbs or solid, continuous herbaceous cover.

Measurement Criteria Herb cover pattern is used to describe the herb layer spatial distribution.

Standard: Herb cover pattern is based on the majority area coverage.

Default:

Permitted Values Herb Cover Pattern Code

1. Single to very few (<4) occurrences of limited extent, circular to irregular shape
2. Single to very few (<4) occurrences of limited extent, linear or elongated shape.
3. Several (>3) sporadic occurrences of limited extent, circular to irregular shape.
4. Several (>3) sporadic occurrences of limited extent, linear or elongated shape.
5. Intimately intermixed units, often with gradational transitions from one to the other.
6. Discontinuous but extensive occurrences, parallel to sub-parallel elongated in shape.
7. Limited continuous occurrence with few inclusions.
8. Continuous occurrence with several inclusions.
9. Continuous occurrence with very few inclusions.

Input Format: #
 Input Example: 3
 Data Origin: input
 Attribute Source: vri

Sequence:	486
Optional:	Y
Format:	number
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: herb_cover_pct Short Name: herb_pct
Alias herb cover pct

Forestry Term: Herb Cover Percentage

Description: Herb cover percent is the percentage of ground area covered by herbaceous cover visible to the photo interpreter. Herb cover percent is analogous to tree and shrub crown closures and is expressed as a percentage of the entire polygon.

Measurement Criteria Herb cover percent provides a direct estimate of herbaceous cover.

Standard: Record herbaceous cover to the nearest percent.

Default:

Permitted Values Integer: 1 to 100

Input Format: ###
Input Example: 10
Data Origin: input
Attribute Source: vri

Sequence:	484
Optional:	Y
Format:	number
Length:	3
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: herb_cover_type Short Name: herb_type
 Alias herb cover type

Forestry Term: Herb Cover Type

Description: This set of attributes describes the portion of herb cover that is not obscured by the vertical projection of the crowns of either trees or shrubs. Herbs are defined as non-woody (vascular) plants, including graminoids (sedges, rushes, grasses), forbs (ferns, club mosses, and horsetails) and some low, woody species and intermediate life forms.

Measurement Criteria

Standard: Measured to the level of resolution that can be photo interpreted for all herbaceous cover types observable in the polygon.

Default:

Permitted Values Codes Description

HE Herb
 A Herb polygon with no distinction between forbs and graminoids

HF Herb - Forbs
 A Herb polygon with forbs greater than 50% of the herb cover.

HG Herb - Graminoids
 A Herb polygon with graminoids greater than 50% of the herb cover.

Input Format: XX
 Input Example: HE
 Data Origin: input
 Attribute Source: vri

Sequence:	482
Optional:	Y
Format:	varchar2
Length:	2
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: input_date Short Name: input_date

Alias input date

Forestry Term: Input Date

Description: The date the forest cover information was entered into the Provincial Data Base.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: MM/DD/YY

Input Example: 10/19/00

Data Origin: input

Attribute Source: vri

Sequence:	22
Optional:	Y
Format:	date
Length:	
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: interpretation_date Short Name: intrp_date

Alias interpretation date

Forestry Term: Interpretation Date

Description: The date on which the polygon estimates were photo interpreted.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: MM/DD/YY

Input Example: 10/19/00

Data Origin: input

Attribute Source: vri

Sequence:	46
Optional:	Y
Format:	date
Length:	
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: interpreted_data_src_cd Short Name: interp_cd
Alias: interpreted data src cd

Forestry Term: Interpreted Data Source Code

Description: The source of the data that contributed to the determination of the classification description. All values taken from Table 3-1, PIP This list of values is similar, but not identical to the FIP DATA SOURCE which will be used to validate the FIP file prior to loading. Non-conforming FIP DATA SOURCE values will be converted to the VEGETATION DATA SOURCE values on load to the Vegetative Cover database.

Measurement Criteria

Standard:

Default:

Permitted Values Codes - Data Sources - Possible Applications
0 - Photo interpretation -

1 - Air call (air observation without 70 mm photography) - species composition

2 - Air call from low - level, fixed base (70 mm photography) - species comp., height

3 - Phase 1 photo sample (pre - 1990) -

4 - Ground call 1 point - age, height

5 - Standard fixed radius sample (pre - 1979) - age, height

6 - Phase 2 or phase 3 sample (pre - 1990) - species, age, height, density, basal area

7 - Silviculture surveys - stocking, survival, free growing, pre - stand tending - species composition, density, SMR, SNR

8 - Ground observation with measurement - age, height

9 - Research plots (e.g. Sx trials, ecological site description) - species, age, height

10 - Valuation cruise plot(s) - basal area, species composition, height

11 - Silviculture treatment record - a record that summarizes the modified stand structure following an activity or treatment such as planting, juvenile spacing, brushing and weeding, conifer release, seed tree control, sanitation spacing, rehabilitation or commercial thinning -

12 - Disturbance - an area recently disturbed by fire, logging, windthrow, or insects that is classified as NSR. Has no source of information other than type and year of disturbance -

13 - Managed stand sample -

14 - Ground call, 2 or more points - age, height, species composition

16 - Vegetation sample - age, height, density, basal area, SMR, SNR

17 - Vegetation ground call - age, height, density, basal area, SMR, SNR

18 - Vegetation air call - species composition, shrub height, shrub %

19 - Natural growth sample - species, age, height

20 - Volume and depletion sample - age, height

22 - Photogrammetrically captured information that is determined or captured using photogrammetric means. An example of this is the determination of photo - measured heights using softcopy technology or parallax bars. - age, height

Input Format: ##
Input Example: 1
Data Origin: input
Attribute Source: vri

Sequence:	8
Optional:	Y
Format:	number
Length:	2
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: interpreter Short Name: interpreter
 Alias: interpreter

Forestry Term: Interpreter

Description: The name of the person who provided the estimates for the data associated with each polygon.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: XXXXXXXX
 Input Example: John Smith
 Data Origin: input
 Attribute Source: vri

Sequence:	30
Optional:	Y
Format:	varchar2
Length:	30
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: inventory_standard_cd Short Name: inv_std_cd
Alias: inventory_standard_cd

Forestry Term: Inventory Standard Code

Description: Code indicating under which inventory standard the data was collected. Values are: "V:" for Vegetation Resources Inventory (VRI), "F" for Forest Inventory Planning (FIP) and "I" for Incomplete (when a full set of VRI attributes is not collected).

Measurement Criteria

Standard:

Default:

Permitted Values V - full VRI;
 F - Forest Inventory Planning (FIP);
 I - Incomplete VRI

Input Format: X
Input Example: V
Data Origin: input
Attribute Source: vri

Sequence:	40
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: label_centre_x Short Name: lbl_ctr_x

Alias label centre x

Forestry Term: Label Centre X

Description: The x co-ordinate of the suggested centre of the label.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	780
Optional:	Y
Format:	number
Length:	38
Decimal Places:	10
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: label_centre_y Short Name: lbl_ctr_y
 Alias label centre y

Forestry Term: Label Centre Y

Description: The y co-ordinate of the suggested centre of the label.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	790
Optional:	Y
Format:	number
Length:	38
Decimal Places:	10
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: label_height Short Name: lbl_ht
 Alias label height

Forestry Term: Label Height

Description: The height of the full label for a 1:15,000 map presentation in meters. It is calculated as 30 times the number of lines in the full label.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	745
Optional:	Y
Format:	number
Length:	38
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: label_width Short Name: lbl_width

Alias label width

Forestry Term: Label Width

Description: The width of the full label for a 1:15,000 map presentation in meters. It is calculated as 18 times the number of characters in the longest line.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	740
Optional:	Y
Format:	number
Length:	38
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: land_cover_class_cd_2 Short Name: land_cd_2

Alias land cover component code

Forestry Term: Land Cover Class Code 2

Description: The Land Cover component Code_2 describes the second most dominate land cover type by percent area occupied within the polygon that contribute to the overall polygon description, but may be too small to be spatially identified. The sub-division of a polygon by a quantified Land Cover Component, allowing non-spatial resolution for modeling of wildlife habitat capability.

Land cover component identifies a type of land cover under the B.C. Land Cover Classification Scheme, to the most detailed level possible (Level 4 for Vegetated; Levels 4 or 5 for Non-Vegetated).

The land cover class code provides a categorization of the polygon to the most detailed land cover description level of the B.C. Land Cover Classification Scheme. This information can be used for the classification of individual polygons and can be summarized for national and international reporting.

Measurement Criteria Enter the appropriate two-letter code for any polygon component that:

- consists of continuous area(s) that are individually greater than or equal to 10% of the polygon area;
- is distinct at levels 4 or 5 of the B.C. Land Cover Classification Scheme; and,
- would otherwise be delineated and classified at approximately twice the map scale.

Standard: Describe up to three land cover components (in decreasing size, by area). Enter the appropriate code under Land Cover Component #1, Land Cover Component #2 and Land Cover Component #3.

If more than three components exist, the remaining percent cover is recorded under "Other Land Cover Component Percent Coverage."

Default:

Permitted Values Land cover component codes - Vegetated
Codes Description

TB Treed Broadleaf

A Treed polygon where 75% or more of the tree basal area, expressed as percentage species composition, consists of broadleaf cover.

TC Treed Coniferous

A Treed polygon where 75% or more of the tree basal area, expressed as percentage species composition, consists of coniferous cover.

TM Treed Mixed

A Treed polygon where neither coniferous nor broadleaf cover individually constitutes at least 75% of the tree basal area, expressed as percentage species composition.

ST Shrub Tall

A Shrub polygon with shrub height of two metres or more.

SL Shrub Low

A Shrub polygon with shrub height less than two metres.

HE Herb

A Herb polygon with no distinction between forbs and graminoids.

HF Herb - Fords

A Herb polygon with forbs greater than 50% of the herb cover.

HG Herb - Graminoids

A Herb polygon with graminoids greater than 50% of the herb cover.

BY Bryoid

A Bryoid polygon with no distinction between mosses and lichens.

BM Bryoid - Moss (bryophytes)

A Bryoid polygon with bryophytes greater than 50% of the bryoid cover.

BL Bryoid - Lichens

A Bryoid polygon with lichens greater than 50% of the bryoid cover.

Land cover component codes - Non-Vegetated**Codes Description****SI Snow / Ice**

Either glacier (which is considered a mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction) or other ice and snow cover that is not part of a glacier.

GL Glacier

A mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction.

PN Snow Cover

Snow or ice that is not part of a glacier, but is found during summer months on the landscape.

RO Rock / Rubble

Bedrock or fragmented rock broken away from bedrock surfaces and moved into its present position by gravity or ice. Extensive deposits are found in and adjacent to alpine areas and are associated with steep rock walls and exposed ridges; canyons and cliff areas also contain these deposits.

BR Bedrock

Unfragmented, consolidated rock contiguous with underlying material.

TA Talus

Rock fragments of any size accumulated on or at the foot of slopes as a result of successive rock falls. This is a type of colluvium.

BI Blockfield

Blocks of rock derived from the underlying bedrock by weathering and/or frost heaving. These have not undergone any significant down slope movement as they occur on level or gently sloping areas.

MZ Rubbly Mine Spoils

Discarded overburden or waste rock, moved to extract ore during a mining operation.

LB Lava Bed

An area where molten rock has flowed from a volcano or fissure and cooled and solidified to form rock.

EL Exposed Land

All other forms of Exposed Land identified by a range of subclasses.

RS River Sediments

Silt, gravel, and sand bars associated with former river channels and present river edges.

ES Exposed Soil

Any exposed soil not covered by other categories, such as areas of recent disturbance that include mud slides, debris torrents, avalanches, or disturbances such as pipeline rights-of-way or cultivated fields where vegetation cover is less than 5%.

LS Pond or Lake Sediments

Exposed sediments related to dried-up lakes or ponds.

RM Reservoir Margin

Land exposed by a drained or fluctuating reservoir. It is found above "normal" water levels and may consist of a range of substrates including gravel, cobbles,

fine sediments, or bedrock.

BE Beach

An area with sorted sediments reworked in recent time by wave action. It may be formed at the edge of fresh or salt water bodies.

LL Landing

A compacted area adjacent to a road used for the purpose of sorting and loading logs.

BU Burned Area

Land showing evidence of recent burning, either natural or prescribed. Vegetation of less than 5% crown cover is present at the time of polygon description.

RZ Road Surface

An area cleared and compacted for transporting goods and services by vehicles. Older roads that are used infrequently or not at all may cease to be classed as Non-Vegetated.

MU Mudflat Sediment

Flat plain-like areas associated with lakes, ponds, rivers, or streams - dominated by fine-textured sediments. They can be associated with freshwater or estuarine sources.

CB Cutbank

Part of a road corridor created upslope of the road surface by excavation into the hillside.

MN Moraine

An area of debris transported and deposited by a glacier.

GP Gravel Pit

An area exposed through the removal of sand and gravel.

TZ Tailings

An area containing the solid waste material produced in the mining and milling of ore.

RN Railway

A roadbed with fixed rails which may contain single or multiple rail lines.

UR Urban

Buildings and associated developments such as roads and parking areas which form an almost continuous covering of the landscape.

AP Airport

A permanent, paved or gravel area, and associated buildings and parking, used by airplanes.

MI Open Pit Mine

An exposed area used to extract ore during a mining operation. This may contain associated buildings and any tailing produced by the mining and milling process.

OT Other

A Non-Vegetated polygon where none of the above categories can be reliably chosen.

Water Cover

LA Lake

A naturally occurring static body of water more than two metres deep in some portion. The boundary for the lake is the natural high water mark.

RE Reservoir

An artificial basin affected by impoundment behind a structure such as a dam, berm, dyke, or wall.

RI River/Stream

A watercourse formed when water flows between continuous, definable banks. Flow may be intermittent or perennial, but does not include ephemeral flow where a channel with no definable banks is present. Gravel bars are part of a stream, while islands within a stream that have definable banks are not.

OC Ocean
A naturally occurring body of water containing salt or generally considered to be salty.

Input Format:
Input Example:
Data Origin:
Attribute Source:

Sequence: Optional: Format: Length: Decimal Places: Null:
--

Use:
Linkage:
Relationship:
Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: land_cover_class_cd_3 Short Name: land_cd_3

Alias: land cover component code

Forestry Term: Land Cover Class Code 3

Description: The Land Cover component Code_3 describes the third most dominate land cover type by percent area occupied within the polygon that contribute to the overall polygon description, but may be too small to be spatially identified. The sub-division of a polygon by a quantified Land Cover Component, allowing non-spatial resolution for modeling of wildlife habitat capability.

Land cover component identifies a type of land cover under the B.C. Land Cover Classification Scheme, to the most detailed level possible (Level 4 for Vegetated; Levels 4 or 5 for Non-Vegetated).

The land cover class code provides a categorization of the polygon to the most detailed land cover description level of the B.C. Land Cover Classification Scheme. This information can be used for the classification of individual polygons and can be summarized for national and international reporting.

Measurement Criteria: Enter the appropriate two-letter code for any polygon component that:

- consists of continuous area(s) that are individually greater than or equal to 10% of the polygon area;
- is distinct at levels 4 or 5 of the B.C. Land Cover Classification Scheme; and,
- would otherwise be delineated and classified at approximately twice the map scale.

Standard: Describe up to three land cover components (in decreasing size, by area). Enter the appropriate code under Land Cover Component #1, Land Cover Component #2 and Land Cover Component #3.

If more than three components exist, the remaining percent cover is recorded under "Other Land Cover Component Percent Coverage."

Default:

Permitted Values: Land cover component codes - Vegetated
Codes Description

TB Treed Broadleaf

A Treed polygon where 75% or more of the tree basal area, expressed as percentage species composition, consists of broadleaf cover.

TC Treed Coniferous

A Treed polygon where 75% or more of the tree basal area, expressed as percentage species composition, consists of coniferous cover.

TM Treed Mixed

A Treed polygon where neither coniferous nor broadleaf cover individually constitutes at least 75% of the tree basal area, expressed as percentage species composition.

ST Shrub Tall

A Shrub polygon with shrub height of two metres or more.

SL Shrub Low

A Shrub polygon with shrub height less than two metres.

HE Herb

A Herb polygon with no distinction between forbs and graminoids.

HF Herb - Forbs

A Herb polygon with forbs greater than 50% of the herb cover.

HG Herb - Graminoids

A Herb polygon with graminoids greater than 50% of the herb cover.

BY Bryoid

A Bryoid polygon with no distinction between mosses and lichens.

BM Bryoid - Moss (bryophytes)

A Bryoid polygon with bryophytes greater than 50% of the bryoid cover.

BL Bryoid - Lichens

A Bryoid polygon with lichens greater than 50% of the bryoid cover.

Land cover component codes - Non-Vegetated**Codes Description****SI Snow / Ice**

Either glacier (which is considered a mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction) or other ice and snow cover that is not part of a glacier.

GL Glacier

A mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction.

PN Snow Cover

Snow or ice that is not part of a glacier, but is found during summer months on the landscape.

RO Rock / Rubble

Bedrock or fragmented rock broken away from bedrock surfaces and moved into its present position by gravity or ice. Extensive deposits are found in and adjacent to alpine areas and are associated with steep rock walls and exposed ridges; canyons and cliff areas also contain these deposits.

BR Bedrock

Unfragmented, consolidated rock contiguous with underlying material.

TA Talus

Rock fragments of any size accumulated on or at the foot of slopes as a result of successive rock falls. This is a type of colluvium.

BI Blockfield

Blocks of rock derived from the underlying bedrock by weathering and/or frost heaving. These have not undergone any significant down slope movement as they occur on level or gently sloping areas.

MZ Rubbly Mine Spoils

Discarded overburden or waste rock, moved to extract ore during a mining operation.

LB Lava Bed

An area where molten rock has flowed from a volcano or fissure and cooled and solidified to form rock.

EL Exposed Land

All other forms of Exposed Land identified by a range of subclasses.

RS River Sediments

Silt, gravel, and sand bars associated with former river channels and present river edges.

ES Exposed Soil

Any exposed soil not covered by other categories, such as areas of recent disturbance that include mud slides, debris torrents, avalanches, or disturbances such as pipeline rights-of-way or cultivated fields where vegetation cover is less than 5%.

LS Pond or Lake Sediments

Exposed sediments related to dried-up lakes or ponds.

RM Reservoir Margin

Land exposed by a drained or fluctuating reservoir. It is found above "normal" water levels and may consist of a range of substrates including gravel, cobbles,

fine sediments, or bedrock.

BE Beach

An area with sorted sediments reworked in recent time by wave action. It may be formed at the edge of fresh or salt water bodies.

LL Landing

A compacted area adjacent to a road used for the purpose of sorting and loading logs.

BU Burned Area

Land showing evidence of recent burning, either natural or prescribed. Vegetation of less than 5% crown cover is present at the time of polygon description.

RZ Road Surface

An area cleared and compacted for transporting goods and services by vehicles. Older roads that are used infrequently or not at all may cease to be classed as Non-Vegetated.

MU Mudflat Sediment

Flat plain-like areas associated with lakes, ponds, rivers, or streams - dominated by fine-textured sediments. They can be associated with freshwater or estuarine sources.

CB Cutbank

Part of a road corridor created upslope of the road surface by excavation into the hillside.

MN Moraine

An area of debris transported and deposited by a glacier.

GP Gravel Pit

An area exposed through the removal of sand and gravel.

TZ Tailings

An area containing the solid waste material produced in the mining and milling of ore.

RN Railway

A roadbed with fixed rails which may contain single or multiple rail lines.

UR Urban

Buildings and associated developments such as roads and parking areas which form an almost continuous covering of the landscape.

AP Airport

A permanent, paved or gravel area, and associated buildings and parking, used by airplanes.

MI Open Pit Mine

An exposed area used to extract ore during a mining operation. This may contain associated buildings and any tailing produced by the mining and milling process.

OT Other

A Non-Vegetated polygon where none of the above categories can be reliably chosen.

Water Cover

LA Lake

A naturally occurring static body of water more than two metres deep in some portion. The boundary for the lake is the natural high water mark.

RE Reservoir

An artificial basin affected by impoundment behind a structure such as a dam, berm, dyke, or wall.

RI River/Stream

A watercourse formed when water flows between continuous, definable banks. Flow may be intermittent or perennial, but does not include ephemeral flow where a channel with no definable banks is present. Gravel bars are part of a stream, while islands within a stream that have definable banks are not.

OC Ocean

A naturally occurring body of water containing salt or generally considered to be salty.

Input Format:
Input Example:
Data Origin:
Attribute Source:

Sequence:
Optional:
Format:
Length:
Decimal Places:
Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: land_cover_class_cd_1 Short Name: land_cd_1

Alias land cover component code

Forestry Term: Land Cover Class Code 1

Description: The Land Cover component Code_1 describes the first most dominate land cover type by percent area occupied within the polygon that contribute to the overall polygon description, but may be too small to be spatially identified. The sub-division of a polygon by a quantified Land Cover Component, allowing non-spatial resolution for modeling of wildlife habitat capability.

Land cover component identifies a type of land cover under the B.C. Land Cover Classification Scheme, to the most detailed level possible (Level 4 for Vegetated; Levels 4 or 5 for Non-Vegetated).

The land cover class code provides a categorization of the polygon to the most detailed land cover description level of the B.C. Land Cover Classification Scheme. This information can be used for the classification of individual polygons and can be summarized for national and international reporting.

Measurement Criteria Enter the appropriate two-letter code for any polygon component that:

- consists of continuous area(s) that are individually greater than or equal to 10% of the polygon area;
- is distinct at levels 4 or 5 of the B.C. Land Cover Classification Scheme; and,
- would otherwise be delineated and classified at approximately twice the map scale.

Standard: Describe up to three land cover components (in decreasing size, by area). Enter the appropriate code under Land Cover Component #1, Land Cover Component #2 and Land Cover Component #3.

If more than three components exist, the remaining percent cover is recorded under "Other Land Cover Component Percent Coverage."

Default:

Permitted Values Land cover component codes - Vegetated
Codes Description

TB Treed Broadleaf

A Treed polygon where 75% or more of the tree basal area, expressed as percentage species composition, consists of broadleaf cover.

TC Treed Coniferous

A Treed polygon where 75% or more of the tree basal area, expressed as percentage species composition, consists of coniferous cover.

TM Treed Mixed

A Treed polygon where neither coniferous nor broadleaf cover individually constitutes at least 75% of the tree basal area, expressed as percentage species composition.

ST Shrub Tall

A Shrub polygon with shrub height of two metres or more.

SL Shrub Low

A Shrub polygon with shrub height less than two metres.

HE Herb

A Herb polygon with no distinction between forbs and graminoids.

HF Herb - Forbs

A Herb polygon with forbs greater than 50% of the herb cover.

HG Herb - Graminoids

A Herb polygon with graminoids greater than 50% of the herb cover.

BY Bryoid

A Bryoid polygon with no distinction between mosses and lichens.

BM Bryoid - Moss (bryophytes)

A Bryoid polygon with bryophytes greater than 50% of the bryoid cover.

BL Bryoid - Lichens

A Bryoid polygon with lichens greater than 50% of the bryoid cover.

Land cover component codes - Non-Vegetated**Codes Description****SI Snow / Ice**

Either glacier (which is considered a mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction) or other ice and snow cover that is not part of a glacier.

GL Glacier

A mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction.

PN Snow Cover

Snow or ice that is not part of a glacier, but is found during summer months on the landscape.

RO Rock / Rubble

Bedrock or fragmented rock broken away from bedrock surfaces and moved into its present position by gravity or ice. Extensive deposits are found in and adjacent to alpine areas and are associated with steep rock walls and exposed ridges; canyons and cliff areas also contain these deposits.

BR Bedrock

Unfragmented, consolidated rock contiguous with underlying material.

TA Talus

Rock fragments of any size accumulated on or at the foot of slopes as a result of successive rock falls. This is a type of colluvium.

BI Blockfield

Blocks of rock derived from the underlying bedrock by weathering and/or frost heaving. These have not undergone any significant down slope movement as they occur on level or gently sloping areas.

MZ Rubbly Mine Spoils

Discarded overburden or waste rock, moved to extract ore during a mining operation.

LB Lava Bed

An area where molten rock has flowed from a volcano or fissure and cooled and solidified to form rock.

EL Exposed Land

All other forms of Exposed Land identified by a range of subclasses.

RS River Sediments

Silt, gravel, and sand bars associated with former river channels and present river edges.

ES Exposed Soil

Any exposed soil not covered by other categories, such as areas of recent disturbance that include mud slides, debris torrents, avalanches, or disturbances such as pipeline rights-of-way or cultivated fields where vegetation cover is less than 5%.

LS Pond or Lake Sediments

Exposed sediments related to dried-up lakes or ponds.

RM Reservoir Margin

Land exposed by a drained or fluctuating reservoir. It is found above "normal" water levels and may consist of a range of substrates including gravel, cobbles,

fine sediments, or bedrock.

BE Beach

An area with sorted sediments reworked in recent time by wave action. It may be formed at the edge of fresh or salt water bodies.

LL Landing

A compacted area adjacent to a road used for the purpose of sorting and loading logs.

BU Burned Area

Land showing evidence of recent burning, either natural or prescribed. Vegetation of less than 5% crown cover is present at the time of polygon description.

RZ Road Surface

An area cleared and compacted for transporting goods and services by vehicles. Older roads that are used infrequently or not at all may cease to be classed as Non-Vegetated.

MU Mudflat Sediment

Flat plain-like areas associated with lakes, ponds, rivers, or streams - dominated by fine-textured sediments. They can be associated with freshwater or estuarine sources.

CB Cutbank

Part of a road corridor created upslope of the road surface by excavation into the hillside.

MN Moraine

An area of debris transported and deposited by a glacier.

GP Gravel Pit

An area exposed through the removal of sand and gravel.

TZ Tailings

An area containing the solid waste material produced in the mining and milling of ore.

RN Railway

A roadbed with fixed rails which may contain single or multiple rail lines.

UR Urban

Buildings and associated developments such as roads and parking areas which form an almost continuous covering of the landscape.

AP Airport

A permanent, paved or gravel area, and associated buildings and parking, used by airplanes.

MI Open Pit Mine

An exposed area used to extract ore during a mining operation. This may contain associated buildings and any tailing produced by the mining and milling process.

OT Other

A Non-Vegetated polygon where none of the above categories can be reliably chosen.

Water Cover

LA Lake

A naturally occurring static body of water more than two metres deep in some portion. The boundary for the lake is the natural high water mark.

RE Reservoir

An artificial basin affected by impoundment behind a structure such as a dam, berm, dyke, or wall.

RI River/Stream

A watercourse formed when water flows between continuous, definable banks. Flow may be intermittent or perennial, but does not include ephemeral flow where a channel with no definable banks is present. Gravel bars are part of a stream, while islands within a stream that have definable banks are not.

OC Ocean
A naturally occurring body of water containing salt or generally considered to be salty.

Input Format:
Input Example:
Data Origin:
Attribute Source:

Sequence: Optional: Format: Length: Decimal Places: Null:
--

Use:
Linkage:
Relationship:
Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: layer_id Short Name: layer_id
Alias layer id

Forestry Term: Layer Identity

Description: The unique business identification of a layer, or horizontal stratum, in a stand. Each layer is normally characterized as a distinct canopy containing a common forest cover structure with timber of similar ages (at least 40 years between layers) and heights (at least 10 meters between layers). Layers are assigned from the tallest layer downward.

Measurement Criteria Each layer is normally characterized as a distinct canopy containing a common forest cover structure with timber of similar ages (at least 40 years between layers) and heights (at least 10 meters between layers).

Standard: Layers are assigned from the tallest layer downward.

Default:

Permitted Values 1
2
3
S
V

Input Format: X or #
Input Example: 1
Data Origin: input
Attribute Source: vri

Sequence:	4
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	N

Use:

Linkage:

Relationship:

Sub Type Links: tree_cover_layer tree_species tree_species_volume vegrpt_polylayer veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: line_1_opening_number Short Name: lbl_opn_no
Alias line 1 opening number

Forestry Term: Label Line 1 Opening Number

Description: Indicates combination of layers and tree class to be summarised for volume.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	705
Optional:	Y
Format:	varchar2
Length:	4
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_1_opening_symbol_cd Short Name: lbl_opn_cd
 Alias: line 1 opening symbol cd

Forestry Term: Label Line 1 Opening Symbol Code

Description: The opening symbol code is represented as one of the following characters: 'X', '|', or '~'. If the opening number is null, line 1 is not populated, so there is no opening symbol. If the adjoining NTS map number is in the form "num num num char num / char", it is an NTS number, the corresponding opening symbol is a hexagon with an 'N' in it, and is represented here by '~'. If the adjoining NTS map number is in the form "num num num char num num num", it is a BCGS number, the corresponding opening symbol is a hexagon with an 'X' in it, and is represented here by '|'. Otherwise the opening symbol is an empty hexagon, and is represented here by 'X'.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:
 Input Example:
 Data Origin:
 Attribute Source:

Sequence:	710
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_2_polygon_id Short Name: lbl_polyid
Alias: line 2 polygon id

Forestry Term: Label Line 2 Polygon Identity

Description: The polygon ID for which this is the label. This is followed by /L (a multi-layered stand) or /S (a separate silviculture description is available in the data base).

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	711
Optional:	Y
Format:	varchar2
Length:	10
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_3_tree_species Short Name: lbl_specis
 Alias: line 3 tree species

Forestry Term: Label Line 3 Tree Species

Description: A list of major species (minor species), ordered by percentage. The species symbols are F (Douglas fir), C (western red cedar), H (hemlock), B (balsam), S (spruce), Sb (black spruce), Yc (yellow cedar), Pw (western white pine), Pa (whitebark pine), Pj (jack pine), L (larch), Ac (cottonwood), D (red alder), Mb (broadleaf maple), E (birch), Al (aspen).

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	712
Optional:	Y
Format:	varchar2
Length:	50
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_4_classes_indexes Short Name: lbl_cls_in
 Alias: line 4 classes indexes

Forestry Term: Label Line 4 Index Classes

Description: Line 4 is made up of 4 numerical characters followed by a hyphen, the site index, a slash, and the estimated site index. The four numerical characters represent projected age class, projected height class, projected stocking class, and crown closure class in that order.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	713
Optional:	Y
Format:	varchar2
Length:	12
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_5_vegetation_cover Short Name: lbl_vegcov
Alias line 5 vegetation cover

Forestry Term: Label Line 5 Vegetation Cover

Description: A listing of the non-vegetated descriptors or the non tree vegetative cover types ordered from most to least common. Possible values in the list are sh (shrub), he (herb), by (bryoid), or the non-vegetative cover codes.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:
Input Example:
Data Origin:
Attribute Source:

Sequence:	714
Optional:	Y
Format:	varchar2
Length:	11
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_6_site_prep_history Short Name: lbl_hist
 Alias: line 6 site prep history

Forestry Term: Label Line 6 Site Preparation History

Description: The site preparation history represented by a list of abbreviations for the techniques used, followed by the years each technique was used. Possible values for the abbreviations are B (broadcast burn) c (chemical), G (grass seeded), H (hand preparation), RB (range management burn), S (spot burn), M (mechanical), MS (mechanical and spot burn), and W (windrow).

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	715
Optional:	Y
Format:	varchar2
Length:	10
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_7_activity_hist_symbol Short Name: lbl_distur
 Alias: line 7 activity hist symbol

Forestry Term: Label Line 7 Activity History Symbol

Description: A symbol representing what techniques were used in the labelled area. The symbol is a circle with 0 to 4 radius lines. Each line represents a technique applied to the labelled area.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:
 Input Example:
 Data Origin:
 Attribute Source:

Sequence:	718
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_7A_stand_tending_history Short Name: lbl_his_sy
 Alias: line 7A stand tending history

Forestry Term: Label Line 7A Stand Tending History

Description: Symbols representing tending history

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	716
Optional:	Y
Format:	varchar2
Length:	39
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_7B_disturbance_history Short Name: lbl_tend
 Alias line 7B disturbance history

Forestry Term: Label Line 7B Disturbance History

Description: The disturbance history described as a list of abbreviations for the techniques along with the years each technique was employed. Possible values are B (wildfire), BE (escaped burn), BG (ground burn), BR (range burn), BW (wildlife burn), D (disease), F (flooding), I (insect), K (fume kill), L (logging), L% (logged with percentage), R (site rehabilitation), S (slide), and W (wind throw).

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	719
Optional:	Y
Format:	varchar2
Length:	40
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: line_8_planting_history Short Name: lbl_plant
 Alias: line 8 planting history

Forestry Term: Label Line 8 Planting History

Description: The planting (or regeneration) history described as a list of years during which artificial plantings was performed.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:
 Input Example:
 Data Origin:
 Attribute Source:

Sequence:	720
Optional:	Y
Format:	varchar2
Length:	80
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type:

Attribute Name: live_stand_volume_125 Short Name: lvoltot_125
 Alias: live stand volume for 12.5 cm

Forestry Term: Live Stand Volume for 12.5cm

Description: This is the total net live volume per hectare of all species determined by percent basal area of the tree layer at the 12.5 cm utilization level. Net volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, net volumes are calculated for rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
 Input Example: 235.65
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: numeric
 Length: 8
 Decimal Places: 3
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_stand_volume_175 Short Name: lvoltot_175
 Alias: live stand volume for 17.5 cm

Forestry Term: Live Stand Volume for 17.5cm

Description: This is the total net live volume per hectare of all species determined by percent basal area of the tree layer at the 17.5 cm utilization level. Net volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, net volumes are calculated for rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
 Input Example: 235.65
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: numeric
 Length: 8
 Decimal Places: 3
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_stand_volume_225 Short Name: lvoltot_225
Alias live stand volume for 22.5 cm

Forestry Term: Live Stand Volume for 22.5cm

Description: This is the total net live volume per hectare of all species determined by percent basal area of the tree layer at the 22.5 cm utilization level. Net volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, net volumes are calculated for rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
Input Example: 235.65
Data Origin: derived
Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp1_125 Short Name: lvolsp1_125

Alias live volume per hectare for leading species at 12.5 cm

Forestry Term: Leading Species Live Volume per Hectare at 12.5 cm

Description: This is the net live volume per hectare of the leading species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the leading species

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp1_175 Short Name: lvolsp1_175

Alias live volume per hectare for leading species at 17.5 cm

Forestry Term: Leading Species Live Volume per Hectare at 17.5 cm

Description: This is the net live volume per hectare of the leading species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the leading species

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp1_225 Short Name: lvolsp1_225

Alias live volume per hectare for leading species at 22.5 cm

Forestry Term: Leading Species Live Volume per Hectare at 22.5 cm

Description: This is the net live volume per hectare of the leading species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the leading species

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp2_125 Short Name: lvolsp2_125

Alias live volume per hectare for second species at 12.5 cm

Forestry Term: Second Species Live Volume per Hectare at 12.5 cm

Description: This is the net live volume per hectare of the second species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp2_175 Short Name: lvolsp2_175

Alias live volume per hectare for second species at 17.5 cm

Forestry Term: Second Species Live Volume per Hectare at 17.5 cm

Description: This is the net live volume per hectare of the second species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp2_225 Short Name: lvolsp2_225

Alias live volume per hectare for second species at 22.5 cm

Forestry Term: Second Species Live Volume per Hectare at 22.5 cm

Description: This is the net live volume per hectare of the second species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp3_125 Short Name: lvolsp3_125

Alias live volume per hectare for third species at 12.5 cm

Forestry Term: Third Species Live Volume per Hectare at 12.5 cm

Description: This is the net live volume per hectare of the third species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp3_175 Short Name: lvolsp3_175

Alias live volume per hectare for third species at 17.5 cm

Forestry Term: Third Species Live Volume per Hectare at 17.5 cm

Description: This is the net live volume per hectare of the third species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp3_225 Short Name: lvolsp3_225

Alias live volume per hectare for third species at 22.5 cm

Forestry Term: Third Species Live Volume per Hectare at 22.5 cm

Description: This is the net live volume per hectare of the third species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp4_125 Short Name: lvolsp4_125

Alias live volume per hectare for fourth species at 12.5 cm

Forestry Term: Fourth Species Live Volume per Hectare at 12.5 cm

Description: This is the net live volume per hectare of the fourth species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp4_175 Short Name: lvolsp4_175
Alias live volume per hectare for fourth species at 17.5 cm

Forestry Term: Fourth Species Live Volume per Hectare at 17.5 cm

Description: This is the net live volume per hectare of the fourth species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
Input Example: 235.65
Data Origin: derived
Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp4_225 Short Name: lvolsp4_225

Alias live volume per hectare for fourth species at 22.5 cm

Forestry Term: Fourth Species Live Volume per Hectare at 22.5 cm

Description: This is the net live volume per hectare of the fourth species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp5_125 Short Name: lvolsp5_125

Alias live volume per hectare for fifth species at 12.5 cm

Forestry Term: Fifth Species Live Volume per Hectare at 12.5 cm

Description: This is the net live volume per hectare of the fifth species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp5_175 Short Name: lvolsp5_175

Alias live volume per hectare for fifth species at 17.5 cm

Forestry Term: Fifth Species Live Volume per Hectare at 17.5 cm

Description: This is the net live volume per hectare of the fifth species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp5_225 Short Name: lvolsp5_225

Alias live volume per hectare for fifth species at 22.5 cm

Forestry Term: Fifth Species Live Volume per Hectare at 22.5 cm

Description: This is the net live volume per hectare of the fifth species determined by percent basal area of the tree layer at the 22.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp6_125 Short Name: lvolsp6_125
 Alias: live volume per hectare for sixth species at 12.5 cm

Forestry Term: Sixth Species Live Volume per Hectare at 12.5 cm

Description: This is the net live volume per hectare of the sixth species determined by percent basal area of the tree layer at the 12.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
 Input Example: 235.65
 Data Origin: derived
 Attribute Source: both

Sequence:
 Optional:
 Format: numeric
 Length: 8
 Decimal Places: 3
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp6_175 Short Name: lvolsp6_175

Alias live volume per hectare for sixth species at 17.5 cm

Forestry Term: Sixth Species Live Volume per Hectare at 17.5 cm

Description: This is the net live volume per hectare of the sixth species determined by percent basal area of the tree layer at the 17.5 cm utilization level. net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: live_vol_per_ha_spp6_225 Short Name: lvolsp6_225

Alias live volume per hectare for sixth species at 22.5 cm

Forestry Term: Sixth Species Live Volume per Hectare at 22.5 cm

Description: This is the net live volume per hectare of the sixth species determined by percent basal area of the tree layer at the 22.5 cm utilization level. Net live volume per hectare is determined as gross volume less decay, waste, and breakage. Depending on the magnitude of the species' decay, waste and breakage, the net live volume for the second species may be lower than volume for other species in the stand. Net live volumes are calculated for Rank 1 layers only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##

Input Example: 235.65

Data Origin: derived

Attribute Source: both

Sequence:

Optional:

Format: numeric

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type: veg_comp_poly

Attribute Name: map_id Short Name: map_id
 Alias map id

Forestry Term: Forest Cover Map Number

Description: Identifies the Vegetation Cover Map corresponding to the VRI file

Measurement Criteria

Standard: 8 character alpha code holding BCGS map number

Default: must have value

Permitted Values The identifier in this case is eight long and is made up of:
 Position 2-4 MAPSHEET GRID NTS or BCGS. Values are 82, 83, 92, 93, 94, 102, 103, 104, 114.
 5 MAPSHEET LETTER BCGS/NTS letter. Values are A - P, and W.
 6-8 MAPSHEET SQUARE BCGS Number or NTS Number and letter. BCGS number values are 1-100, and NTS number values are 1-16 with NTS letter values A-H , and W.
 9 MAPSHEET QUAD an identifier for 3' x 6' (1:10,000 scale) mapsheets.
 e.g.. 082G002 - 6' x 12' minute map sheet

Input Format: XXXXXXXX
 Input Example: 093J034
 Data Origin: input
 Attribute Source: vri

Sequence:	
Optional:	
Format:	varchar2
Length:	7
Decimal Places:	
Null:	N

Use: Identifies the mapsheet containing the corresponding Forest Cover Map.

Linkage:

Relationship:

Sub Type Links: non_vegetative_cover tree_cover_layer tree_layer_history_link tree_species tree_species_volume
 vegetative_cover veg_data_set_version veg_label veg_vegetation_cover_polygon veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: modifying_process Short Name: mod_proces
Alias: modifying process

Forestry Term: Modifying Process

Description: A natural mechanism of weathering, erosion and soil material deposition that result in the modification of surficial materials and landforms. Used for terrain classification, site classification, soil condition and identification of potential hazards such as avalanches, slope instability and flooding.

Measurement Criteria: Only active modifying processes are to be assigned. A process is considered active if there is evidence of current or recent occurrence or likely future occurrence.

Standard: The code is recorded for the prevalent modifying process within the polygon on the basis of percent area coverage.

Default:

Permitted Values: A Avalanching Slopes modified by the rapid downslope movement of snow and ice and by the deposition of rock debris, surficial material and vegetation debris transported by snow avalanches. Sites usually contain avalanche chutes and run out zones but may also be affected by ice falling from glaciers.

B River channelling Erosion and channel formation by the flow of water within clearly defined banks.

F Mass movements Down slope movement of cohesive or non-cohesive surficial material and/or bedrock by creeping, sliding, flowing or falling. This includes rock and debris slides, soil slumps and talus slopes.

N None of these descriptions apply; no modifying processes are observed in the polygon.

U Flooding Areas subject to periodic (possibly seasonal) inundation with subsequent deposition of soil particles. Commonly applied to ephemeral lakes.

V Gully erosion Modification of unconsolidated or consolidated surfaces by processes such as running water and snow avalanching that result in the formation of parallel or sub-parallel long, narrow ravines. Singular gullies are not generally included in this class.

Input Format: X
Input Example: F
Data Origin: input
Attribute Source: vri

Sequence:	31
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: Used for terrain classification, site classification, soil condition and identification of potential hazards such as avalanches, slope instability and flooding.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: non_forest_descriptor Short Name: nfor_desc
Alias non forest descriptor

Forestry Term: Non Forest Descriptor

Description: A classification code indicating that the forest cover type is not currently forested, but is capable of supporting commercial forests. This is a FIP classification based attribute only, and is retained for the purposes of business transition from FIP to Vegetation Inventory

Measurement Criteria The Non-Forest Descriptor indicates that the forest cover polygon is potentially productive, but is not currently supporting commercial forests.

Standard: 5 character alpha holding the abbreviation for Non-Forest descriptor.

Default:

Permitted Values NCBR- Non-commercial brush
NC- Non-Commercial
NSR - Not sufficiently restocked
NP - Non-Productive
NTA - No typing available

Input Format: XXXXX
Input Example: NCBR
Data Origin: input
Attribute Source: fip

Sequence:
Optional:
Format: varchar2
Length: 5
Decimal Places:
Null: Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes: Used to define land that is not currently forested but is capable of supporting commercial forest. It is also used to determine potential areas for silviculture treatment and to determine the net land base for Timber Supply Analyses.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_productive_cd Short Name: np_code
Alias non productive cd

Forestry Term: Non Productive Code

Description: A unique numeric code that references the classes or type of non-productive areas or land that is incapable of supporting commercial forests. This is a FIP classification based attribute only, and is retained for the purposes of business transition from FIP to Vegetation Inventory. There is no expectation that this attribute would be updated or created under Vegetation Inventory classification practise.

Measurement Criteria

Standard: 2 Character numeric code designating non-productive type code.

Default:

- Permitted Values
- 01 ICE - Icefield
 - 02 A - Alpine
 - 03 R - Rock
 - 06 GR - Gravel Pit
 - 07 SAND - Sand
 - 09 CL - Clay Bank
 - 10 AF - Alpine Forest (with species etc.)
 - 11 NPBR - Non-Productive Brush
 - 12 NP - Non-Productive
 - 12 NP - Non-Productive Forest (with species etc.)
 - 13 NPBU - Non-Productive Burn
 - 15 L - Lake
 - 16 TIDE - Tidal Flat
 - 18 G - Gravel Bar
 - 25 RIV - River
 - 26 MUD - Mud Flat
 - 35 S - (for input) Swamp (completed file)
 - 42 C - Clearing
 - 50 U - Roads
 - 54 U - Urban
 - 60 P - Hayfield
 - 62 M - Meadow
 - 63 OR - Open Range
 - 64 NA - Non-Applicable (salt water)

Input Format: ##
Input Example: 03
Data Origin: input
Attribute Source: vri

Sequence:
Optional:
Format: number
Length: 2
Decimal Places:
Null: Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_productive_descriptor_cd Short Name: np_desc
Alias: non productive descriptor cd

Forestry Term: Non Productive Descriptor Code

Description: A unique code that references the classes or type of non-productive areas. This is a FIP classification based attribute only, and is retained for the purposes of business transition from FIP to Vegetation Inventory. There is no expectation that this attribute would be updated or created under Vegetation Inventory classification practise.

Measurement Criteria Used to provide area summaries and statistics for various classes of non-productive areas.

Standard: 5 character alpha code holding the abbreviation of the non-productive descriptor.

Default:

Permitted Values NTA No Typing Available
ICE Icefield
A Alpine
R Rock
GR Gravel Pit
SAND Sand
CL Clay Bank
AF Alpine Forest (with Species etc.)
NPBR Non-Productive Brush
NP Non-Productive
NP Non-Productive Forest (with species etc.)
NPBU Non-Productive Burn
L Lake
TIDE Tidal Flat
G Gravel Bar
RIV River
MUD Mud Flat
S Swamp (muskeg)
C Clearing
U Roads
U Urban
P Hayfield
M Meadow
OR Open Range
NA Non-Applicable (salt water)

Input Format: XXXXX
Input Example: ICE
Data Origin: input
Attribute Source: vri

Sequence:	20
Optional:	Y
Format:	varchar2
Length:	5
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: A
Non-Productive descriptor does not imply that the land is unproductive for other valuable resources, such as wildlife, fisheries, recreation, etc.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_veg_cover_pattern_1 Short Name: nveg_cov_1
 Alias: non vegetation cover pattern 1

Forestry Term: Non Vegetation Cover Pattern One

Description: Non-vegetated cover pattern_1 describes the spatial distribution of the predominate non-vegetated cover type based on percent area covered within the polygon. Each non-vegetated cover type indicated must have a non-vegetated cover pattern assigned. Non-vegetated cover pattern is used to describe non-vegetated cover spatial distribution.

Examples include roads within vegetated polygons or lakes within non-vegetated polygons.

Measurement Criteria: There are no constraints on the number of non-vegetated cover types to be identified.

Standard:

Default:

Permitted Values: 1 to 9

1. Single to very few (<4) occurrences of limited extent, circular to irregular shape.
2. Single to very few (<4) occurrences of limited extent, linear or elongated shape.
3. Several (>3) sporadic occurrences of limited extent, circular to irregular shape.
4. Several (>3) sporadic occurrences of limited extent, linear or elongated shape.
5. Intimately intermixed units, often with gradational transitions from one to the other.
6. Discontinuous but extensive occurrences, parallel to sub-parallel elongated in shape.
7. Limited continuous occurrence with few inclusions.
8. Continuous occurrence with several inclusions.
9. Continuous occurrence with very few inclusions.

Input Format: #
 Input Example: 5
 Data Origin: input
 Attribute Source: vri

Sequence:
 Optional:
 Format: number
 Length: 1
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_veg_cover_pattern_2 Short Name: nveg_cov_2
 Alias: non vegetation cover pattern 2

Forestry Term: Non Vegetation Cover Pattern 2

Description: Non-vegetated cover pattern_2 describes the spatial distribution of the predominate non-vegetated cover type based on percent area covered within the polygon. Each non-vegetated cover type indicated must have a non-vegetated cover pattern assigned. Non-vegetated cover pattern is used to describe non-vegetated cover spatial distribution.

Examples include roads within vegetated polygons or lakes within non-vegetated polygons.

Measurement Criteria: There are no constraints on the number of non-vegetated cover types to be identified.

Standard:

Default:

Permitted Values: 1 to 9

1. Single to very few (<4) occurrences of limited extent, circular to irregular shape.
2. Single to very few (<4) occurrences of limited extent, linear or elongated shape.
3. Several (>3) sporadic occurrences of limited extent, circular to irregular shape.
4. Several (>3) sporadic occurrences of limited extent, linear or elongated shape.
5. Intimately intermixed units, often with gradational transitions from one to the other.
6. Discontinuous but extensive occurrences, parallel to sub-parallel elongated in shape.
7. Limited continuous occurrence with few inclusions.
8. Continuous occurrence with several inclusions.
9. Continuous occurrence with very few inclusions.

Input Format: #
 Input Example: 4
 Data Origin: input
 Attribute Source: vri

Sequence:
 Optional:
 Format: number
 Length: 1
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_veg_cover_pattern_3 Short Name: nveg_cov_3

Alias non vegetation cover pattern 3

Forestry Term: Non Vegetation Cover Pattern 3

Description: Non-vegetated cover pattern_3 describes the spatial distribution of the predominate non-vegetated cover type based on percent area covered within the polygon. Each non-vegetated cover type indicated must have a non-vegetated cover pattern assigned. Non-vegetated cover pattern is used to describe non-vegetated cover spatial distribution.

Examples include roads within vegetated polygons or lakes within non-vegetated polygons.

Measurement Criteria There are no constraints on the number of non-vegetated cover types to be identified.

Standard:

Default:

Permitted Values 1 to 9

1. Single to very few (<4) occurrences of limited extent, circular to irregular shape.
2. Single to very few (<4) occurrences of limited extent, linear or elongated shape.
3. Several (>3) sporadic occurrences of limited extent, circular to irregular shape.
4. Several (>3) sporadic occurrences of limited extent, linear or elongated shape.
5. Intimately intermixed units, often with gradational transitions from one to the other.
6. Discontinuous but extensive occurrences, parallel to sub-parallel elongated in shape.
7. Limited continuous occurrence with few inclusions.
8. Continuous occurrence with several inclusions.
9. Continuous occurrence with very few inclusions.

Input Format: #
 Input Example: 3
 Data Origin: input
 Attribute Source: vri

Sequence:
 Optional:
 Format: number
 Length: 1
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_veg_cover_pct_1 Short Name: nveg_pct_1
 Alias non vegetation cover percentage 1

Forestry Term: Non Vegetation Cover Percentage One

Description: The area the predominate non-vegetated portion covers expressed as a percentage of the entire polygons area. Non-vegetated cover percent indicates the percentage of the polygon area that is occupied by a non-vegetated cover.

Measurement Criteria Estimate non-vegetated cover percent, based on the percentage of ground area of the polygon covered by the non-vegetated cover type. Enter the cover percent estimate on the same line of the attribute form as the cover type. Enter the non-vegetated cover percent for each non-vegetated cover type identified in the polygon. There are no constraints to the number of non-vegetated cover types that can be identified.

Standard:

Default:

Permitted Values 1 to 100

Input Format: ###
 Input Example: 4
 Data Origin: input
 Attribute Source: vri

Sequence:
 Optional:
 Format: number
 Length: 3
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_veg_cover_pct_2 Short Name: nveg_pct_2

Alias non vegetation cover percentage 2

Forestry Term: Non Vegetation Cover Percentage Two

Description: The area the second most prevalent non-vegetated portion covers expressed as a percentage of the entire polygons area. Non-vegetated cover percent indicates the percentage of the polygon area that is occupied by a non-vegetated cover.

Measurement Criteria Estimate non-vegetated cover percent, based on the percentage of ground area of the polygon covered by the non-vegetated cover type. Enter the cover percent estimate on the same line of the attribute form as the cover type. Enter the non-vegetated cover percent for each non-vegetated cover type identified in the polygon. There are no constraints to the number of non-vegetated cover types that can be identified.

Standard:

Default:

Permitted Values 1 to 100

Input Format: ###

Input Example: 4

Data Origin: input

Attribute Source: vri

Sequence:

Optional:

Format: number

Length: 3

Decimal Places:

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_veg_cover_pct_3 Short Name: nveg_pct_3

Alias non vegetation cover percentage 3

Forestry Term: Non Vegetation Cover Percentage 3

Description: The area the third most prevalent non-vegetated portion covers expressed as a percentage of the entire polygons area. Non-vegetated cover percent indicates the percentage of the polygon area that is occupied by a non-vegetated cover.

Measurement Criteria Estimate non-vegetated cover percent, based on the percentage of ground area of the polygon covered by the non-vegetated cover type. Enter the cover percent estimate on the same line of the attribute form as the cover type. Enter the non-vegetated cover percent for each non-vegetated cover type identified in the polygon. There are no constraints to the number of non-vegetated cover types that can be identified.

Standard:

Default:

Permitted Values 1 to 100

Input Format: ###

Input Example: 4

Data Origin: input

Attribute Source: vri

Sequence:

Optional:

Format: number

Length: 3

Decimal Places:

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_veg_cover_type_1 Short Name: nveg_typ_1
 Alias non vegetation cover type 1

Forestry Term: Non Vegetation Cover Type One

Description: Non-vegetated cover type_1 is the designation for the predominate observable non-vegetated land cover within the polygon. Non-vegetated cover types provide detailed reporting for non-vegetated land cover.

Measurement Criteria Enter the appropriate code to the level of resolution that can be photo interpreted for all non-vegetated cover types observable within the polygon. If more than one non-vegetated cover type is identified, use additional rows on the attribute form.

Standard:

Default:

Permitted Values	CODES DESCRIPTION
	Land Cover
	GL Glacier A mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction.
	PN Snow Cover Snow or ice that is not part of a glacier but is found during summer months on the landscape.
	BR Bedrock Unfragmented, consolidated rock, contiguous with underlying material.
	TA Talus Rock fragments of any size accumulated on or at the foot of slopes as a result of successive rock falls. This is a type of colluvium.
	BI Blockfield Blocks of rock derived from the underlying bedrock by weathering and / or frost heaving. These have not undergone any significant down slope movement as they occur on level or gently sloping areas.
	MZ Rubbly Mine Spoils Discarded overburden or waste rock, moved to extract ore during mining.
	LB Lava Bed An area where molten rock has flowed from a volcano or fissure and cooled and solidified to form rock.
	RS River Sediments Silt, gravel and sand bars associated with former river channels and present river edges.
	ES Exposed Soil Any exposed soil not covered by other categories such as areas of recent disturbance that include mud slides, debris torrents, avalanches, or disturbances such as pipeline rights-of-way where vegetation cover is less than five percent.
	LS Pond or Lake Sediments Exposed sediments related to dried lakes or ponds.
	RM Reservoir Margin Land exposed by a drained or fluctuating reservoir. This is found above "normal" water levels and may consist of a range of substrates including gravel, cobbles, fine sediments, or bedrock.
	BE Beach

An area with sorted sediments reworked in recent time by wave action which may be formed at the edge of fresh or salt water bodies

LL Landing

A compacted area adjacent to a road used for the purpose of sorting and loading logs.

BU Burned Area

Land showing evidence of recent burning, either natural or prescribed. Vegetation of less than 5% crown cover is present at the time of polygon description.

RZ Road Surface

An area cleared and compacted for transporting goods and services by vehicles. Older roads that are used infrequently or not at all may cease to be non-vegetated.

MU Mudflat Sediment

Flat plain-like areas associated with lakes, ponds, rivers or streams - dominated by fine textured sediments. They can be associated with freshwater or estuarine sources.

CB Cutbank

Part of a road corridor created up slope of the road surface created by excavation into the hillside.

MN Moraine

An area of debris transported and deposited by a glacier.

GP Gravel Pit

An area exposed through the removal of sand and gravel.

TZ Tailings

An area containing the solid waste material produced in the mining and milling of ore.

RN Railway

A roadbed with fixed rails which may contain single or multiple rail lines.

UR Urban

Buildings and associated developments such as roads and parking areas which form an almost continuous covering of the landscape.

AP Airport

A permanent, paved or gravel area, and associated buildings and parking, used by airplanes.

MI Open Pit Mine

An exposed area used to extract ore during a mining operation. This may contain associated buildings and any tailing produced by the mining and milling process.

OT Other

A non-vegetated polygon where none of the above categories can be reliably chosen.

Water Cover

LA Lake

A naturally occurring static body of water more than two metres deep in some portion. The boundary for the lake is the natural high water mark.

RE Reservoir

An artificial basin affected by impoundment behind a man made structure such as a dam, berm, dyke, or wall.

RI River/Stream

A water course formed when water flows between continuous, definable banks. Flow may be intermittent or perennial but does not include ephemeral flow where a channel with no definable banks is present.

Gravel bars are part of a stream while islands within a stream that have definable banks are not.

OC Ocean
A naturally occurring body of water containing salt or generally considered to be salty.

Input Format: XX
Input Example: RE
Data Origin: input
Attribute Source: vri

Sequence:	
Optional:	
Format:	varchar2
Length:	2
Decimal Places:	
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_veg_cover_type_2 Short Name: nveg_typ_2

Alias non vegetation cover type 2

Forestry Term: Non Vegetation Cover Type Two

Description: Non-vegetated cover type_2 is the designation for the second most prevalent observable non-vegetated land cover within the polygon. Non-vegetated cover types provide detailed reporting for non-vegetated land cover.

Measurement Criteria Enter the appropriate code to the level of resolution that can be photo interpreted for all non-vegetated cover types observable within the polygon. If more than one non-vegetated cover type is identified, use additional rows on the attribute form.

Standard:

Default:

Permitted Values CODES DESCRIPTION
 Land Cover

GL Glacier
 A mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction.

PN Snow Cover
 Snow or ice that is not part of a glacier but is found during summer months on the landscape.

BR Bedrock
 Unfragmented, consolidated rock, contiguous with underlying material.

TA Talus
 Rock fragments of any size accumulated on or at the foot of slopes as a result of successive rock falls. This is a type of colluvium.

BI Blockfield
 Blocks of rock derived from the underlying bedrock by weathering and / or frost heaving. These have not undergone any significant down slope movement as they occur on level or gently sloping areas.

MZ Rubbly Mine Spoils
 Discarded overburden or waste rock, moved to extract ore during mining.

LB Lava Bed
 An area where molten rock has flowed from a volcano or fissure and cooled and solidified to form rock.

RS River Sediments
 Silt, gravel and sand bars associated with former river channels and present river edges.

ES Exposed Soil
 Any exposed soil not covered by other categories such as areas of recent disturbance that include mud slides, debris torrents, avalanches, or disturbances such as pipeline rights-of-way where vegetation cover is less than five percent.

LS Pond or Lake Sediments
 Exposed sediments related to dried lakes or ponds.

RM Reservoir Margin
 Land exposed by a drained or fluctuating reservoir. This is found above "normal" water levels and may consist of a range of substrates including gravel, cobbles, fine sediments, or bedrock.

BE Beach

An area with sorted sediments reworked in recent time by wave action which may be formed at the edge of fresh or salt water bodies

LL Landing

A compacted area adjacent to a road used for the purpose of sorting and loading logs.

BU Burned Area

Land showing evidence of recent burning, either natural or prescribed. Vegetation of less than 5% crown cover is present at the time of polygon description.

RZ Road Surface

An area cleared and compacted for transporting goods and services by vehicles. Older roads that are used infrequently or not at all may cease to be non-vegetated.

MU Mudflat Sediment

Flat plain-like areas associated with lakes, ponds, rivers or streams - dominated by fine textured sediments. They can be associated with freshwater or estuarine sources.

CB Cutbank

Part of a road corridor created up slope of the road surface created by excavation into the hillside.

MN Moraine

An area of debris transported and deposited by a glacier.

GP Gravel Pit

An area exposed through the removal of sand and gravel.

TZ Tailings

An area containing the solid waste material produced in the mining and milling of ore.

RN Railway

A roadbed with fixed rails which may contain single or multiple rail lines.

UR Urban

Buildings and associated developments such as roads and parking areas which form an almost continuous covering of the landscape.

AP Airport

A permanent, paved or gravel area, and associated buildings and parking, used by airplanes.

MI Open Pit Mine

An exposed area used to extract ore during a mining operation. This may contain associated buildings and any tailing produced by the mining and milling process.

OT Other

A non-vegetated polygon where none of the above categories can be reliably chosen.

Water Cover**LA Lake**

A naturally occurring static body of water more than two metres deep in some portion. The boundary for the lake is the natural high water mark.

RE Reservoir

An artificial basin affected by impoundment behind a man made structure such as a dam, berm, dyke, or wall.

RI River/Stream

A water course formed when water flows between continuous, definable banks. Flow may be intermittent or perennial but does not include

ephemeral flow where a channel with no definable banks is present.
Gravel bars are part of a stream while islands within a stream that have definable banks are not.

OC Ocean
A naturally occurring body of water containing salt or generally considered to be salty.

Input Format: XX
Input Example: RN
Data Origin: input
Attribute Source: vri

Sequence:	
Optional:	
Format:	varchar2
Length:	2
Decimal Places:	
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: non_veg_cover_type_3 Short Name: nveg_typ_3

Alias non vegetation cover type 3

Forestry Term: Non Vegetation Cover Type Three

Description: Non-vegetated cover type_3 is the designation for the third most prevalent observable non-vegetated land cover within the polygon. Non-vegetated cover types provide detailed reporting for non-vegetated land cover.

Measurement Criteria Enter the appropriate code to the level of resolution that can be photo interpreted for all non-vegetated cover types observable within the polygon. If more than one non-vegetated cover type is identified, use additional rows on the attribute form.

Standard:

Default:

Permitted Values CODES DESCRIPTION
Land Cover

GL Glacier
A mass of perennial snow and ice with definite lateral limits, typically flowing in a particular direction.

PN Snow Cover
Snow or ice that is not part of a glacier but is found during summer months on the landscape.

BR Bedrock
Unfragmented, consolidated rock, contiguous with underlying material.

TA Talus
Rock fragments of any size accumulated on or at the foot of slopes as a result of successive rock falls. This is a type of colluvium.

BI Blockfield
Blocks of rock derived from the underlying bedrock by weathering and / or frost heaving. These have not undergone any significant down slope movement as they occur on level or gently sloping areas.

MZ Rubbly Mine Spoils
Discarded overburden or waste rock, moved to extract ore during mining.

LB Lava Bed
An area where molten rock has flowed from a volcano or fissure and cooled and solidified to form rock.

RS River Sediments
Silt, gravel and sand bars associated with former river channels and present river edges.

ES Exposed Soil
Any exposed soil not covered by other categories such as areas of recent disturbance that include mud slides, debris torrents, avalanches, or disturbances such as pipeline rights-of-way where vegetation cover is less than five percent.

LS Pond or Lake Sediments
Exposed sediments related to dried lakes or ponds.

RM Reservoir Margin
Land exposed by a drained or fluctuating reservoir. This is found above "normal" water levels and may consist of a range of substrates including gravel, cobbles, fine sediments, or bedrock.

BE Beach

An area with sorted sediments reworked in recent time by wave action which may be formed at the edge of fresh or salt water bodies

LL Landing

A compacted area adjacent to a road used for the purpose of sorting and loading logs.

BU Burned Area

Land showing evidence of recent burning, either natural or prescribed. Vegetation of less than 5% crown cover is present at the time of polygon description.

RZ Road Surface

An area cleared and compacted for transporting goods and services by vehicles. Older roads that are used infrequently or not at all may cease to be non-vegetated.

MU Mudflat Sediment

Flat plain-like areas associated with lakes, ponds, rivers or streams - dominated by fine textured sediments. They can be associated with freshwater or estuarine sources.

CB Cutbank

Part of a road corridor created up slope of the road surface created by excavation into the hillside.

MN Moraine

An area of debris transported and deposited by a glacier.

GP Gravel Pit

An area exposed through the removal of sand and gravel.

TZ Tailings

An area containing the solid waste material produced in the mining and milling of ore.

RN Railway

A roadbed with fixed rails which may contain single or multiple rail lines.

UR Urban

Buildings and associated developments such as roads and parking areas which form an almost continuous covering of the landscape.

AP Airport

A permanent, paved or gravel area, and associated buildings and parking, used by airplanes.

MI Open Pit Mine

An exposed area used to extract ore during a mining operation. This may contain associated buildings and any tailing produced by the mining and milling process.

OT Other

A non-vegetated polygon where none of the above categories can be reliably chosen.

Water Cover**LA Lake**

A naturally occurring static body of water more than two metres deep in some portion. The boundary for the lake is the natural high water mark.

RE Reservoir

An artificial basin affected by impoundment behind a man made structure such as a dam, berm, dyke, or wall.

RI River/Stream

A water course formed when water flows between continuous, definable banks. Flow may be intermittent or perennial but does not include

ephemeral flow where a channel with no definable banks is present.
Gravel bars are part of a stream while islands within a stream that have definable banks are not.

OC Ocean
A naturally occurring body of water containing salt or generally considered to be salty.

Input Format: XX
Input Example: BU
Data Origin: input
Attribute Source: vri

Sequence:	
Optional:	
Format:	varchar2
Length:	2
Decimal Places:	
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: objectid Short Name: objectid

Alias object id

Forestry Term: Object Identity

Description:

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:

Optional:

Format: number

Length:

Decimal Places:

Null:

Use:

Linkage:

Relationship:

Sub Type Links: veg_vegetation_cover_polygon veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: open_id Short Name: open_id
Alias open id

Forestry Term: Opening Indicator

Description: System generated value uniquely identifying the opening in the MOFR Forest Tenures Administration (FTA) and the RESULTS and Forest Inventory data sets

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:

Optional:

Format:

Length:

Decimal Places:

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: opening_ind Short Name: open_ind
Alias opening_ind

Forestry Term: Opening Indicator

Description: Indicates whether or not the polygon represents a silviculture opening

Measurement Criteria

Standard:

Default:

Permitted Values Y - Yes; N - No

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	
Optional:	
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name:	opening_number	Short Name:	open_num
Alias	opening number		

Forestry Term: Opening Number

Description: A unique number assigned to each opening in the forest caused by a disturbance (e.g. fire, logging, etc.) for which there will be management activities

Measurement Criteria

Standard:

Default:

Permitted Values Numeric value 0 to 9999

Input Format: #####

Input Example: 465

Data Origin: input

Attribute Source: fip

Sequence:

Optional:

Format:	varchar2
Length:	4
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: Identifies Silviculture Opening Number and provides a cross-reference to the Silviculture Data Base(s).

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: opening_source Short Name: open_src

Alias opening source

Forestry Term: Opening Source

Description: Defines whether the opening came from ISIS or MLSIS This field is not populated in the current data model.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	
Optional:	
Format:	varchar2
Length:	5
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: org_unit_code Short Name: orgunit_cd
Alias: organisation unit code

Forestry Term: Organisation Unit Code

Description: Forest District Codes, 3 charcter description

Measurement Criteria

Standard:

Default:

Permitted Values	org. number	org unit code	org unit name
	=====	=====	=====
	1833	RNI	Northern Interior Forest Region
	46	DFN	Fort Nelson Forest District
	34	DJA	Fort St. James Forest District
	32	DKM	Kalum Forest District
	38	DMK	Mackenzie Forest District
	1823	DND	Nadina Forest District
	1825	DPC	Peace Forest District
	18	DPG	Prince George Forest District
	1824	DSS	Skeena Stikine Forest District
	DVA	DVA	Vanderhoof Forest District
	1834	RSI	Southern Interior Forest Region
	56	DMH	100 Mile House Forest District
	1830	DAB	Arrow Boundary Forest District
	1828	DCS	Cascades Forest District
	1826	DCC	Central Cariboo Forest District
	58	DCH	Chilcotin Forest District
	1620	DCO	Columbia Forest District
	1827	DHW	Headwaters Forest District
	21	DKA	Kamloops Forest District
	60	DKL	Kootena Lake Forest District
	1829	DOS	Okanagan Shuswap Forest District
	50	DQU	Quesnel Forest District
	1831	DRM	Rocky Mountain Forest District
	1835	RCO	Coast Forest Region
	43	DCR	Campbell River Forest District
	15	DCK	Chilliwack Forest District
	36	DNC	North Coast Forest District
	1832	DNI	North Island - Central Coast Forest District
	48	DQC	Queen Charlotte Islands Forest District
	1619	DSI	South Island Forest District
	23	DSQ	Squamish Forest District
	27	DSC	Sunshine Coast Forest District

Input Format: XXX
Input Example: DFN
Data Origin: derived
Attribute Source: vri

Sequence:
Optional:
Format: varchar2
Length: 3
Decimal Places:
Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: printable_ind Short Name: printable

Alias printable ind

Forestry Term: Printable

Description: "Y" means print the label. "N" means do not print the label.

Measurement Criteria

Standard:

Default:

Permitted Values Y - Yes; N - No

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	735
Optional:	N
Format:	varchar2
Length:	1
Decimal Places:	
Null:	N

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_layer

Attribute Name: proj_age_1 Short Name: proj_age_1
 Alias: projected age 1

Forestry Term: Projected Age for Leading Species

Description: Projected age for leading species

Measurement Criteria

Standard:

Default:

Permitted Values greater or equal than 1

Input Format: ###
 Input Example: 120
 Data Origin: input / projected
 Attribute Source: vri

Sequence:
 Optional:
 Format: number
 Length: 4
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: proj_age_2 Short Name: proj_age_2
 Alias: projected age 2

Forestry Term: Projected Age for Second Species

Description: Projected age for secondary species

Measurement Criteria

Standard:

Default:

Permitted Values greater or greater than 1

Input Format: ###
 Input Example: 120
 Data Origin: input / projected
 Attribute Source: vri

Sequence:
 Optional:
 Format: number
 Length: 4
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: proj_age_class_cd_1 Short Name: p_age_cas1
 Alias projected age class code 1

Forestry Term: Projected age class code stand age 1

Description: The age class projected to the current year for species 1.

Measurement Criteria

Standard:

Default: 0

Permitted Values
 1 Stand age 1 to 20 years
 2 Stand age 21 to 40 years
 3 Stand age 41 to 60 years
 4 Stand age 61 to 80 years
 5 Stand age 81 to 100 years
 6 Stand age 101 to 120 years
 7 Stand age 121 to 140 years
 8 Stand age 141 to 250 years
 9 Stand age 251 + years

Input Format: #
 Input Example: 2
 Data Origin: derived
 Attribute Source:

Sequence:
 Optional:
 Format: number
 Length: 4
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type:

Attribute Name: proj_age_class_cd_2 Short Name: p_age_cas2
 Alias projected age class code 2

Forestry Term: Projected age class code stand age 2

Description: The age class projected to the current year for species 1.

Measurement Criteria

Standard:

Default: 0

Permitted Values
 1 Stand age 1 to 20 years
 2 Stand age 21 to 40 years
 3 Stand age 41 to 60 years
 4 Stand age 61 to 80 years
 5 Stand age 81 to 100 years
 6 Stand age 101 to 120 years
 7 Stand age 121 to 140 years
 8 Stand age 141 to 250 years
 9 Stand age 251 + years

Input Format: #
 Input Example: 2
 Data Origin: derived
 Attribute Source:

Sequence:
 Optional:
 Format: number
 Length: 4
 Decimal Places:
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type: veg_comp_layer

Attribute Name: proj_height_1 Short Name: proj_ht_1
 Alias: projected height 1

Forestry Term: Projected Height for Leading Species

Description: Projected height for leading species

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ##.#
 Input Example: 29.4
 Data Origin: input / projected
 Attribute Source: vri

Sequence:	
Optional:	
Format:	number
Length:	5
Decimal Places:	1
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: proj_height_2 Short Name: proj_ht_2
 Alias: projected height 2

Forestry Term: Projected Height for Second Species

Description: projected height for second species

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ##.#
 Input Example: 29.4
 Data Origin: input / projected
 Attribute Source: vri

Sequence:
 Optional:
 Format: number
 Length: 5
 Decimal Places: 1
 Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: proj_height_class_cd_1 Short Name: p_ht_cas1
 Alias projected height class code 1

Forestry Term: Projected height class code 1
 Description: The height class projected to the current year for species 1.

Measurement Criteria

Standard:

Default: 0
 Permitted Values 0 0 m 1 > 0.0 - 10.4 m
 1 0.1 - 10.4 m
 2 10.5 - 19.4 m
 3 19.5 - 28.4 m
 4 28.5 - 37.4 m
 5 37.5 - 46.4 m
 6 46.5 - 55.4 m
 7 55.5 - 64.4 m
 8 64.5 +

Input Format: #
 Input Example: 2
 Data Origin: derived
 Attribute Source:

Sequence:
 Optional:
 Format: number
 Length: 4
 Decimal Places:
 Null:

Use:
 Linkage:
 Relationship:
 Sub Type Links:

Notes:
 Tips and Hints:
 Reference:

Sub Type:

Attribute Name: proj_height_class_cd_2 Short Name: p_ht_cas2
 Alias: projected height class code 2

Forestry Term: Projected height class code 2
 Description: The height class projected to the current year for species 2.

Measurement Criteria

Standard:

Default: 0
 Permitted Values: 0 0 m 1 > 0.0 - 10.4 m
 1 0.1 - 10.4 m
 2 10.5 - 19.4 m
 3 19.5 - 28.4 m
 4 28.5 - 37.4 m
 5 37.5 - 46.4 m
 6 46.5 - 55.4 m
 7 55.5 - 64.4 m
 8 64.5 +

Input Format: #
 Input Example: 2
 Data Origin: derived
 Attribute Source:

Sequence:
 Optional:
 Format: number
 Length: 4
 Decimal Places:
 Null:

Use:
 Linkage:
 Relationship:
 Sub Type Links:

Notes:
 Tips and Hints:
 Reference:

Sub Type: veg_comp_poly

Attribute Name: project Short Name: project_id
Alias project

Forestry Term: Project

Description: The business assigned name of the project. The name typically reflects a Timber Supply Area, an initiating Agency, or a land area.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: XXXXXXXXXXXXXXXXXXXX
Input Example: Cassiar Soft Copy Retrofit
Data Origin: input
Attribute Source: vri

Sequence:	47
Optional:	
Format:	varchar2
Length:	30
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: projected_date Short Name: proj_date
 Alias: projected date

Forestry Term: Projected Date

Description: The date to which time dependent stand information is projected. Used to determine the date to which time dependent variables in the stand have been projected. Attributes that are projected to a future date include: -Age, Age Class, Height, Height Class, Type Identity, Stocking Class, All maps within a project area should be projected to the same date.

Measurement Criteria: Attributes that are projected to a future date include: Age, Age Class, Height, Height Class, Type Identity, Stocking Class, etc.

Standard: 8 character numeric code holding year, month and day

Default:

Permitted Values

Input Format: MMDDYYYY

Input Example: 06012008

Data Origin: derived

Attribute Source: both

Sequence:
 Optional:
 Format: date
 Length:
 Decimal Places:
 Null: Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: quad_diam_125 Short Name: q_diam_125

Alias: quadratic diameter for 12.5 cm

Forestry Term: Quadratic Diamter at 12.5 cm

Description: The quadratic mean stand diameter (breast height), at the projection date, based on the 12.5 cm utilization level. Calculated for Rank 1 stands only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: #####.###

Input Example:

Data Origin: derived

Attribute Source:

Sequence:

Optional:

Format: number

Length: 8

Decimal Places: 3

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: quad_diam_175 Short Name: q_diam_175

Alias quadratic diameter for 17.5 cm

Forestry Term: Quadratic Diamter at 17.5 cm

Description: The quadratic mean stand diameter (breast height), at the projection date, based on the 17.5 cm utilization level. Calculated for Rank 1 stands only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: #####.###

Input Example:

Data Origin: derived

Attribute Source:

Sequence:	
Optional:	
Format:	number
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: quad_diam_225 Short Name: q_diam_225
 Alias quadratic diameter for 22.5 cm

Forestry Term: Quadratic Diamter at 22.5 cm

Description: The quadratic mean stand diameter (breast height), at the projection date, based on the 22.5 cm utilization level. Calculated for Rank 1 stands only.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: #####.###

Input Example:

Data Origin: derived

Attribute Source:

Sequence:	
Optional:	
Format:	number
Length:	8
Decimal Places:	3
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: reference_date Short Name: ref_date
 Alias: reference date

Forestry Term: Reference Date

Description: The date of the source data on which the interpretation is based. Known as the 'Reference Year' in the VIF file. In the VRI this is calculated from the year of the photo or source survey that was used to generate the VRI attribute.

Measurement Criteria

Standard: 2 character numeric code indicating year for which the attributes are most reliable.

Default: 53 to present year.

Permitted Values: must have value

Input Format: MMDDYYYY

Input Example: 06012008

Data Origin: derived

Attribute Source: both

Sequence:	48
Optional:	Y
Format:	date
Length:	
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: reference_year Short Name: ref_yr_id
Alias: reference year

Forestry Term: Reference Year

Description: The year of the source data on which the interpretation is based. Known as the 'Reference Year' in the VIF file. In the VRI this is calculated from the year of the photo or source survey that was used to generate the VRI attribute.

Measurement Criteria

Standard: 4 character numeric code indicating year for which the attributes are most reliable

Default: must have value

Permitted Values: 1953 >

Input Format: xxxx
Input Example: 2008
Data Origin: derived
Attribute Source: both

Sequence:	
Optional:	
Format:	number
Length:	4
Decimal Places:	
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: shrub_cover_pattern Short Name: shrb_patt
 Alias: shrub cover pattern

Forestry Term: Shrub Cover Pattern

Description: Shrub cover pattern is a code that describes the spatial distribution of the shrubs within the polygon. Shrub cover pattern is used to describe the shrub layer spatial distribution. Examples include clumps of shrubs on rocky patches or individual shrubs or solid, continuous shrub cover.

Measurement Criteria Shrub cover pattern is used to describe the shrub layer spatial distribution.

Standard: Shrub cover pattern is based on the majority area coverage.

Default:

Permitted Values Shrub Cover Pattern Code

1. Single to very few (<4) occurrences of limited extent, circular to irregular shape.
2. Single to very few (<4) occurrences of limited extent, linear or elongated shape.
3. Several (>3) sporadic occurrences of limited extent, circular to irregular shape.
4. Several (>3) sporadic occurrences of limited extent, linear or elongated shape.
5. Intimately intermixed units, often with gradational transitions from one to the other.
6. Discontinuous but extensive occurrences, parallel to sub-parallel elongated in shape.
7. Limited continuous occurrence with few inclusions.
8. Continuous occurrence with several inclusions.
9. Continuous occurrence with very few inclusions.

Input Format: #
 Input Example: 3
 Data Origin: input
 Attribute Source: vri

Sequence:	480
Optional:	Y
Format:	number
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: shrub_crown_closure Short Name: shrb_cc

Alias shrub crown closure

Forestry Term: Shrub Crown Closure

Description: Shrub crown closure is the percentage of ground area covered by the vertically projected crowns of the shrub cover visible to the photo interpreter. Shrub crown closure is expressed as a percentage of the entire polygon.

Measurement Criteria Estimate crown closure for all shrub species based on the percentage of ground area covered by the vertically projected crowns of shrubs. Only those shrubs not otherwise obscured by tree crown cover are recorded.

Standard: Shrub crown closure is expressed as a percentage of the entire polygon.

Default:

Permitted Values 1 to 100

Input Format: ###

Input Example: 25

Data Origin: input

Attribute Source: vri

Sequence: 470
 Optional: Y
 Format: number
 Length: 3
 Decimal Places:
 Null: Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: Shrub crown closure provides a direct estimate of crown closure that is not adjusted by the Ground Sampling.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: shrub_height Short Name: shrb_ht

Alias shrub height

Forestry Term: Shrub Height

Description: The average height of the shrubs contained in the polygon as interpreted from medium scale photography.

Measurement Criteria Estimate the average height in metres (weighted by crown closure) of all shrubs within the polygon that are not obscured by tree crown cover.

Standard: Shrub crown closure is expressed as a percentage of the entire polygon.

Default:

Permitted Values

Input Format: ##.#

Input Example: 1.5

Data Origin: input

Attribute Source: vri

Sequence:	460
Optional:	Y
Format:	number
Length:	4
Decimal Places:	1
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: When multiplied by shrub cover, an index of shrub volume is obtained that indicates available browse.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: site_index Short Name: site_index

Alias site index

Forestry Term: Site Index

Description: Site index is an estimate of site productivity for tree growth (height in metres at breast height age of 50 years). The mean height of the dominant and codominant trees will attain at a base index age used for the purposes of estimating forestsite growth capability. The site index is based on a normalized set of coefficients calibrated to reflect the range of heights for a given tree species.

Measurement Criteria Estimated site index may be based on the direct application of conventional site index curves, or it may be estimated from other data sources.

Standard: Estimated site index is recorded to the nearest one metre.
4 character numeric value for site index in metres at 50 bha (Breast Height Age)

Default:

Permitted Values

Input Format: ##.#

Input Example: 15.0

Data Origin: derived

Attribute Source: both

Sequence:	
Optional:	
Format:	number
Length:	4
Decimal Places:	1
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes: The direct site index value may be determined from the dominant and codominant trees. Used extensively in Timber Supply Analyses and Local Resource Use Plans (LRUPs) for determining the Contributing Land Base and assigning net-downs.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: site_position_meso Short Name: site_meso

Alias site position meso

Forestry Term: Site Position Meso

Description: A code denoting the relative position of the sampling site within a catchment area with the intent to be consistent within the scale of topography affecting surface water flow. The vertical difference is usually between 3 and 300m, and the surface area generally exceeds 0.5 has in size. Also known as slope position or meso site position.

Measurement Criteria he scale of vertical distance for site position meso is usually between 3 m and 300 m.

Standard: A code is recorded for each polygon for the prevalent site position meso of the polygon on the basis of percent area coverage.

Default:

Permitted Values Codes Description

- C Crest
The generally convex uppermost portion of a hill (meso scale). It is usually convex in all directions and generally has no distinct aspect. The term "crest" may also be applied to a ridge.
- U Upper slope
The generally convex, upper portion of the slope of a hill (meso scale) immediately below the crest. It has a convex surface profile with a specific aspect.
- M Middle slope
The area of the slope of a hill between the upper and lower slope, where the slope profile is not generally concave or convex. It has a straight or somewhat sigmoid surface profile with a specific aspect.
- L Lower slope
The area toward the base of the slope of the hill. It generally has a concave surface profile with a specific aspect.
- T Toe
The area differentiated from the lower slope by an abrupt decrease in slope gradient. It is often characterized by seepage.
- D Depression
Any area that is concave in all directions. It is generally at the foot of a meso scale hill or in a generally level area.
- F Flat (Level)
Any level area not immediately adjacent to a meso scale hill (or toe). The surface profile is generally horizontal with no significant aspect.

Input Format: X
 Input Example: M
 Data Origin: input
 Attribute Source: vri

Sequence:	32
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: Site position meso is one of the key attributes for site series identification. Identification of soil moisture regime, using environmental properties, is done with reference to categories of site position meso.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: small_label Short Name: sm_label

Alias small Label

Forestry Term: Small Label

Description: The two-line (or format 3) version of the label. This label contains, at most, 2 lines build from the line 1 and 2 attributes. A back slash represents a carriage return.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format:

Input Example:

Data Origin:

Attribute Source:

Sequence:	725
Optional:	N
Format:	varchar2
Length:	100
Decimal Places:	
Null:	N

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory

Sub Type: veg_comp_poly

Attribute Name: soil_moisture_regime _2 Short Name: soil_mst_2

Alias soil moisture regime 2

Forestry Term: Soil Moisture Regime 2

Description: The average amount of soil water annually available for evapotranspiration by vascular plants averaged over many years within the second most dominate cover type. Soil Moisture Regime is an intrepretive attribute for estimation of site potential and site series classification.

Measurement Criteria

Standard:

Default:

Permitted Values Soil moisture regime classes

- Codes SMR
- 0 very xeric
- 1 xeric
- 2 subxeric
- 3 submesic
- 4 mesic
- 5 subhygric
- 6 hygric
- 7 subhydric
- 8 hydric

Input Format: #

Input Example: 4

Data Origin: input

Attribute Source:

Sequence:

Optional:

Format: varchar2

Length: 10

Decimal Places:

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name:	soil_moisture_regime _3	Short Name:	soil_mst_3
Alias	soil moisture regime 3		

Forestry Term: Soil Moisture Regime 3

Description: The average amount of soil water annually available for evapotranspiration by vascular plants averaged over many years within the second most dominate cover type. Soil Moisture Regime is an intrepretive attribute for estimation of site potential and site series classification.

Measurement Criteria

Standard:

Default:

Permitted Values Soil moisture regime classes

- Codes SMR
- 0 very xeric
- 1 xeric
- 2 subxeric
- 3 submesic
- 4 mesic
- 5 subhygric
- 6 hygric
- 7 subhydric
- 8 hydric

Input Format: #

Input Example: 2

Data Origin: input

Attribute Source:

Sequence:

Optional:

Format: varchar2

Length: 10

Decimal Places:

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name:	soil_moisture_regime_1	Short Name:	soil_mst_1
Alias	soil moisture regime 1		

Forestry Term: Soil Moisture Regime 1

Description: The average amount of soil water annually available for evapotranspiration by vascular plants averaged over many years within the predominate cover type. Soil Moisture Regime is an interpretive attribute for estimation of site potential and site series classification.

Measurement Criteria

Standard:

Default:

Permitted Values Soil moisture regime classes

- Codes SMR
- 0 very xeric
- 1 xeric
- 2 subxeric
- 3 submesic
- 4 mesic
- 5 subhygric
- 6 hygric
- 7 subhydric
- 8 hydric

Input Format: #

Input Example: 3

Data Origin: input

Attribute Source:

Sequence:

Optional:

Format: varchar2

Length: 10

Decimal Places:

Null:

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: soil_nutrient_regime Short Name: soil_nutr

Alias: soil nutrient regime

Forestry Term: Soil Nutrient Regime

Description: A code to denote, on a relative scale, the available nutrient supply for plant growth. The soil's nutrient regime (trophotope) integrates many environmental and biotic parameters which, in combination, determine the actual amounts of available nutrients.

Measurement Criteria: Dominant polygon SMR is derived from the largest land cover component by area. If the first two or more land cover components are equal in percent area, the first land cover component indicated will be used to derive the dominant SMR.

Standard: The code is recorded for the dominant SNR of the polygon on the basis of percent area coverage.

Default:

Permitted Values: A Very poor
 B Poor
 C Medium
 D Rich
 E Very rich
 F Ultra rich (saline, excess accumulations of variety of salts).

Input Format: X
 Input Example: D
 Data Origin: input
 Attribute Source: vri

Sequence:	34
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: Soil nutrient regime is an interpretive attribute which, together with soil moisture regime, is used to assist in site series identification.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: special_cruise_number Short Name: cruise_no
Alias: special cruise number

Forestry Term: Special Cruise Number

Description: The numeric code of the Public Sustained Yield Unit(s) (PSYU) that fall within the forest cover polygon. PSYUs are areas of land, usually a natural topographic unit determined by drainage areas. Includes PSYUs, Tree Farm Licences (TFL), Tree Farms (TF), Major Parks and Ecological Reserves, Woodlot licences, and miscellaneous areas.

Measurement Criteria Includes PSYUs, Tree Farm Licences (TFL), Tree Farms (TF), Major Parks and Ecological Reserves, Woodlot licences, and miscellaneous areas.

Standard: 4 character numeric code which references the respective PSYU, TFL, etc.

Default:

Permitted Values 9999 - areas outside PSYU

Input Format: ####
Input Example: 131
Data Origin: derived
Attribute Source: both

Sequence:
Optional:
Format: number
Length: 4
Decimal Places:
Null: Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_poly

Attribute Name: special_cruise_number_cd Short Name: cruise_cd
Alias: special cruise number cd

Forestry Term: Special Cruise Number Code

Description: The numbers of the Public Sustained Yield Unit (PSYU) Block(s) that fall within the forest cover polygon. PSYU Blocks are subdivisions of a PSYU, and indicate the presence of a sub-unit survey (i.e. 1:10,000 scale inventory).

Measurement Criteria

Standard: 1 character alpha code indicating a sub-unit survey

Default:

Permitted Values <blank> No sub-unit survey, Salt Water
9 Sub-unit exist

Input Format: #
Input Example: 9
Data Origin: derived
Attribute Source: both

Sequence:
Optional:
Format: varchar2
Length: 1
Decimal Places:
Null: Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_cd_1 Short Name: spec_cd_1
 Alias: species cd 1

Forestry Term: Species Composition Code - Leading Species

Description: A code describing the leading commercial species or brush species in the layer. The species with the highest percent composition (e.g. gross volume or, if a very young stand, the relative number of stems per hectare) is identified as the leading commercial species. Species must be above a specified diameter to be recognized in the species composition of the layer. Leading species are described in terms of Genus, Species and Subspecies. There are currently 27 commercial tree species and five genus values recognized in the Province. The code may also be used to describe brush species in cases where the Non-Productive Descriptor is NPBR or the Non-Forest Descriptor is NCBR.

Measurement Criteria: The species with the highest percent composition is identified as the leading commercial species. Leading species are described in terms of Genus, Species and Subspecies.

Standard: 3 character alpha code indicating commercial species.

Default: blank

Permitted Values: <blank> No species recorded

Commercial Species

AC	Balsam poplar	Populus balsamifera Black
	Cottonwood	Populus balsamifera
AT	Aspen	Populus tremuloides
B	True fir	Abies spp.
BL	Alpine fir	Abies lasiocarpa
BA	Amabilis fir	Abies amabilis
BG	Grand fir	Abies grandis
CW	Western red cedar	Thuja plicata
DR	Red Alder	Alnus rubra
E	Birch	Betula spp.
EP	Common paper birch	Betula papyrifera
EA	Alaska paper birch	Betula neoalaskana
FD	Douglas fir	Pseudotsuga menziesii
H	Hemlocks	Tsuga spp.
HW	Western hemlock	Tsuga heterophylla
HM	Mountain hemlock	Tsuga mertensiana
L	Larch	Larix spp.
LA	Alpine larch	Larix lyalli
LT	Tamarack	Larix laricina
LW	Western larch	Larix occidentalis
MB	Broadleaf maple	Acer macrophyllum
PF	Limber pine	Pinus flexilis
PL	Lodgepole pine	Pinus contorta
PW	Western white pine	Pinus monticola
PA	Whitebark pine	Pinus albicalis
PY	Yellow pine	Pinus ponderosa
PJ	Jack pine	Pinus banksiana
S	Spruce	Picea spp.
SB	Black spruce	Picea mariana
SE	Engelmann spruce	Picea engelmannii
SS	Sitka spruce	Picea sitchensis
SW	White spruce	Picea glauca
YC	Yellow cedar	Chamaecyparis nootkatensis

Brush Species

DM	Mountain alder	Alnus incana
R	Arbutus	Arbutus menziesii
EW	Water birch	Betula occidentalis

Cedar Thuja C

western redcedar Thuja plicata Cw

Cypress Chamaecyparis Y

yellow-cedar *C. nootkatensis* Yc

Douglas-fir *Pseudotsuga* F
 Douglas-fir *P. menziesii* Fd
 coastal Douglas-fir *P. menziesii* var. *menziesii* Fdc
 interior Douglas-fir *P. menziesii* var. *glauca* Fdi

Fir (Balsam) *Abies* B
 amabilis fir *A. amabilis* Ba
 grand fir *A. grandis* Bg
 subalpine fir *A. lasiocarpa* BI

Hemlock *Tsuga* H
 mountain hemlock *T. mertensiana* Hm
 western hemlock *T. heterophylla* Hw
 mountain x western hemlock hybrid *T. mertensiana* x *heterophylla* Hxm

Juniper *Juniperus* J
 Rocky Mtn. juniper *J. scopulorum* Jr

Larch *Larix* L
 alpine larch *L. lyallii* La
 tamarack *L. laricina* Lt
 western larch *L. occidentalis* Lw

Pine *Pinus* P
 jack pine *P. banksiana* Pj
 limber pine *P. flexilis* Pf
 lodgepole pine *P. contorta* Pl
 lodgepole pine *P. contorta* var. *latifolia* Pli
 lodgepole x jack pine hybrid *P. x murraybanksiana* Pjx
 ponderosa pine *P. ponderosa* Py
 shore pine *P. contorta* var. *contorta* Plc
 western white pine *P. monticola* Pw
 whitebark pine *P. albicaulis* Pa

Spruce *Picea* S
 black spruce *P. mariana* Sb
 Engelmann spruce *P. engelmannii* Se
 Sitka spruce *P. sitchensis* Ss
 white spruce *P. glauca* Sw
 spruce hybrid *Picea* cross Sx
 Engelmann x white *P. engelmannii* x *glauca* Sxw
 Sitka x white *P. x lutzii* Sxl
 Sitka x unknown hybrid *P. sitchensis* x ? Sxs

Yew *Taxus* T
 western yew *Taxus brevifolia* Tw

NATIVE HARDWOODS

Alder *Alnus* D
 red alder *A. rubra* Dr

Apple *Malus* U
 Pacific crab apple *Malus fusca* Up

Aspen, Cottonwood or Poplar *Populus* A
 poplar *P. balsamifera* Ac
 balsam poplar *P. b. ssp. balsamifera* Acb
 black cottonwood *P. b. ssp. trichocarpa* Act
 hybrid poplars *P. spp.* Ax
 trembling aspen *P. tremuloides* At

Arbutus *Arbutus* R
 Arbutus *Arbutus menziesii* Ra

Birch *Betula* E
 Alaska paper birch *B. neoalaskana* Ea

Alaska x paper birch hybrid B. x winteri Exp
 paper birch B. papyrifera Ep
 water birch B. occidentalis Ew

Cascara Rhamnus K
 cascara R. purshiana Kc

Cherry Prunus V
 bitter cherry P. emarginata Vb
 choke cherry P. virginiana Vv
 pin cherry P. pensylvanica Vp

Dogwood Cornus G
 Pacific dogwood Cornus nuttallii Gp

Maple Acer M
 bigleaf maple A. macrophyllum Mb
 vine maple A. circinatum Mv

Oak Quercus Q
 Garry oak Q. garryana Qg

Willow Salix spp. W
 Bebb's willow S. bebbiana Wb
 Pacific willow S. lucida Wp
 peachleaf willow S. amygdaloides Wa
 pussy willow S. discolor Wd
 Scouler's willow S. scouleriana Ws
 Sitka willow S. sitchensis Wt

UNKNOWNNS

Unknown X
 Unknown conifer Xc
 Unknown hardwood Xh

OTHERS

Other tree, not on list Z
 Other conifer Zc
 Other hardwood Zh

EXOTICS

Apple Malus U
 apple Malus pumila Ua

Aspen, Cottonwood or Poplar Populus A
 *southern cottonwood P. deltoides Ad

Birch Betula E
 European birch B. pendula Ee
 silver birch B. pubescens Es
 *yellow birch B. alleghaniensis Ey

Cherry Prunus V
 sweet cherry P. avium Vs

Cypress Chamaecyparis Y
 *Port Orford-cedar C. lawsoniana Yp

Fir (Balsam) Abies B
 *balsam fir A. balsamea Bb
 noble fir A. procera Bp
 *Shasta red fir A. magnifica var. shastensis Bm
 *white fir A. concolor Bc

Larch Larix L
 *Dahurian larch *L. gmelinii* Ld

Maple Acer M
 box elder *A. negundo* Me
 *Norway maple *A. platanoides* Mn
 *Sycamore maple *A. pseudoplatanus* Ms

Oak Quercus Q
 *English oak *Q. robur* Qe
 *white oak *Q. alba* Qw

Other exotics
 *incense-cedar *Calocedrus decurrens* Oa
 *giant sequoia *Sequoiadendron giganteum* Ob
 *coast redwood *Sequoia sempervirens* Oc
 European mountain-ash *Sorbus aucuparia* Od
 Siberian elm *Ulmus pumila* Oe
 common pear *Pyrus communis* Of
 Oregon ash *Fraxinus latifolia* Og
 *white ash *Fraxinus americana* Oh
 *shagbark hickory *Carya ovata* Oi

Pine Pinus P
 *Monterey pine *P. radiata* Pm
 *red pine *P. resinosa* Pr
 *sugar pine *P. lambertiana* Ps

Spruce Picea S
 *Norway spruce *P. abies* Sn

Changes to Version 4 of B.C. Ministry of Forests Tree Code List

1. Exotic species added to list: a) to provide codes for database purposes (requested by Resources Inventory Branch); and b) to accommodate inventories being conducted near areas of settlement that may encounter escaped or naturalized exotics.
2. Hybrids that cannot be easily distinguished or are of doubtful existence were deleted from list. Operational option is to go to upper level generic code if suspected hybrids without codes are encountered; common hybrids still have codes. Deletions include: Sxe (*Picea engelmannii* x *sitchensis*), Sxb (*Picea glauca* x *mariana*), Sxx (*Picea glauca* x *engelmannii* x *sitchensis*), and Exw (*Betula occidentalis* x *papyrifera*).
3. Dm (*Alnus tenuifolia*) deleted as it is not known to exceed 10 m in height and is most often multi-stemmed.

Changes to Version 4.1 of B.C. Ministry of Forests Tree Code List

Sn (*Picea abies*) Norway Spruce was added. Note that some printed copies of version 4.0 had Norway spruce included but our master version required the update hence the minor version upgrade.

Changes to Version 4.2 of B.C. Ministry of Forests Tree Code List

Four exotic species requiring codes for database purposes were added: yellow birch (Ey), white ash (Oh), shagbark hickory (Oi), and white oak (Qw).

Changes to Version 4.3 of B.C. Ministry of Forests Tree Code List

One exotic species requiring a code for database purposes was added: Dahurian larch (Ld).

Species Composition Code - Leading Species

Input Format: XXX
Input Example: PL
Data Origin: input
Attribute Source: both

Sequence:	
Optional:	
Format:	varchar2
Length:	4
Decimal Places:	
Null:	Y

Use: The species code is used in determining: species composition, stand volumes, stand decay, waste and breakage, net-downs in Timber Supply Analyses, site index, etc.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_cd_2 Short Name: spec_cd_2
 Alias: species cd 2

Forestry Term: Species Composition Code - Second Species

Description: A code describing the leading commercial species or brush species in the layer. The species with the highest percent composition (e.g. gross volume or, if a very young stand, the relative number of stems per hectare) is identified as the leading commercial species. Species must be above a specified diameter to be recognized in the species composition of the layer. Leading species are described in terms of Genus, Species and Subspecies. There are currently 27 commercial tree species and five genus values recognized in the Province. The code may also be used to describe brush species in cases where the Non-Productive Descriptor is NPBR or the Non-Forest Descriptor is NCBR.

Measurement Criteria

Standard: 3 character alpha code indicating commercial species.

Default: blank

Permitted Values <blank> No species recorded

Commercial Species

AC	Balsam poplar	Populus balsamifera Black
	Cottonwood	Populus balsamifera
AT	Aspen	Populus tremuloides
B	True fir	Abies spp.
BL	Alpine fir	Abies lasiocarpa
BA	Amabilis fir	Abies amabilis
BG	Grand fir	Abies grandis
CW	Western red cedar	Thuja plicata
DR	Red Alder	Alnus rubra
E	Birch	Betula spp.
EP	Common paper birch	Betula papyrifera
EA	Alaska paper birch	Betula neoalaskana
FD	Douglas fir	Pseudotsuga menziesii
H	Hemlocks	Tsuga spp.
HW	Western hemlock	Tsuga heterophylla
HM	Mountain hemlock	Tsuga mertensiana
L	Larch	Larix spp.
LA	Alpine larch	Larix lyalli
LT	Tamarack	Larix laricina
LW	Western larch	Larix occidentalis
MB	Broadleaf maple	Acer macrophyllum
PF	Limber pine	Pinus flexilis
PL	Lodgepole pine	Pinus contorta
PW	Western white pine	Pinus monticola
PA	Whitebark pine	Pinus albicalis
PY	Yellow pine	Pinus ponderosa
PJ	Jack pine	Pinus banksiana
S	Spruce	Picea spp.
SB	Black spruce	Picea mariana
SE	Engelmann spruce	Picea engelmannii
SS	Sitka spruce	Picea sitchensis
SW	White spruce	Picea glauca
YC	Yellow cedar	Chamaecyparis nootkatensis

Brush Species

DM	Mountain alder	Alnus incana
R	Arbutus	Arbutus menziesii
EW	Water birch	Betula occidentalis

Cedar Thuja C

western redcedar Thuja plicata Cw

Cypress Chamaecyparis Y

yellow-cedar C. nootkatensis Yc

Douglas-fir *Pseudotsuga* F
 Douglas-fir *P. menziesii* Fd
 coastal Douglas-fir *P. menziesii* var. *menziesii* Fdc
 interior Douglas-fir *P. menziesii* var. *glauca* Fdi

Fir (Balsam) *Abies* B
 amabilis fir *A. amabilis* Ba
 grand fir *A. grandis* Bg
 subalpine fir *A. lasiocarpa* BI

Hemlock *Tsuga* H
 mountain hemlock *T. mertensiana* Hm
 western hemlock *T. heterophylla* Hw
 mountain x western hemlock hybrid *T. mertensiana* x *heterophylla* Hxm

Juniper *Juniperus* J
 Rocky Mtn. juniper *J. scopulorum* Jr

Larch *Larix* L
 alpine larch *L. lyallii* La
 tamarack *L. laricina* Lt
 western larch *L. occidentalis* Lw

Pine *Pinus* P
 jack pine *P. banksiana* Pj
 limber pine *P. flexilis* Pf
 lodgepole pine *P. contorta* Pl
 lodgepole pine *P. contorta* var. *latifolia* Pli
 lodgepole x jack pine hybrid *P. x murraybanksiana* Pxi
 ponderosa pine *P. ponderosa* Py
 shore pine *P. contorta* var. *contorta* Plc
 western white pine *P. monticola* Pw
 whitebark pine *P. albicaulis* Pa

Spruce *Picea* S
 black spruce *P. mariana* Sb
 Engelmann spruce *P. engelmannii* Se
 Sitka spruce *P. sitchensis* Ss
 white spruce *P. glauca* Sw
 spruce hybrid *Picea* cross Sx
 Engelmann x white *P. engelmannii* x *glauca* Sxw
 Sitka x white *P. x lutzii* Sxl
 Sitka x unknown hybrid *P. sitchensis* x ? Sxs

Yew *Taxus* T
 western yew *Taxus brevifolia* Tw

NATIVE HARDWOODS

Alder *Alnus* D
 red alder *A. rubra* Dr

Apple *Malus* U
 Pacific crab apple *Malus fusca* Up

Aspen, Cottonwood or Poplar *Populus* A
 poplar *P. balsamifera* Ac
 balsam poplar *P. b. ssp. balsamifera* Acb
 black cottonwood *P. b. ssp. trichocarpa* Act
 hybrid poplars *P. spp.* Ax
 trembling aspen *P. tremuloides* At

Arbutus *Arbutus* R
 Arbutus *Arbutus menziesii* Ra

Birch *Betula* E
 Alaska paper birch *B. neoalaskana* Ea
 Alaska x paper birch hybrid *B. x winteri* Exp

paper birch *B. papyrifera* Ep
water birch *B. occidentalis* Ew

Cascara *Rhamnus* K
cascara *R. purshiana* Kc

Cherry *Prunus* V
bitter cherry *P. emarginata* Vb
choke cherry *P. virginiana* Vv
pin cherry *P. pensylvanica* Vp

Dogwood *Cornus* G
Pacific dogwood *Cornus nuttallii* Gp

Maple *Acer* M
bigleaf maple *A. macrophyllum* Mb
vine maple *A. circinatum* Mv

Oak *Quercus* Q
Garry oak *Q. garryana* Qg

Willow *Salix* spp. W
Bebb's willow *S. bebbiana* Wb
Pacific willow *S. lucida* Wp
peachleaf willow *S. amygdaloides* Wa
pussy willow *S. discolor* Wd
Scouler's willow *S. scouleriana* Ws
Sitka willow *S. sitchensis* Wt

UNKNOWNNS

Unknown X
Unknown conifer Xc
Unknown hardwood Xh

OTHERS

Other tree, not on list Z
Other conifer Zc
Other hardwood Zh

EXOTICS

Apple *Malus* U
apple *Malus pumila* Ua

Aspen, Cottonwood or Poplar *Populus* A
*southern cottonwood *P. deltoides* Ad

Birch *Betula* E
European birch *B. pendula* Ee
silver birch *B. pubescens* Es
*yellow birch *B. alleghaniensis* Ey

Cherry *Prunus* V
sweet cherry *P. avium* Vs

Cypress *Chamaecyparis* Y
*Port Orford-cedar *C. lawsoniana* Yp

Fir (Balsam) *Abies* B
*balsam fir *A. balsamea* Bb
noble fir *A. procera* Bp
*Shasta red fir *A. magnifica* var. *shastensis* Bm
*white fir *A. concolor* Bc

Larch *Larix* L

*Dahurian larch *L. gmelinii* Ld

Maple Acer M

box elder *A. negundo* Me*Norway maple *A. platanoides* Mn*Sycamore maple *A. pseudoplatanus* Ms

Oak Quercus Q

*English oak *Q. robur* Qe*white oak *Q. alba* Qw

Other exotics

*incense-cedar *Calocedrus decurrens* Oa*giant sequoia *Sequoiadendron giganteum* Ob*coast redwood *Sequoia sempervirens* OcEuropean mountain-ash *Sorbus aucuparia* OdSiberian elm *Ulmus pumila* Oecommon pear *Pyrus communis* OfOregon ash *Fraxinus latifolia* Og*white ash *Fraxinus americana* Oh*shagbark hickory *Carya ovata* Oi

Pine Pinus P

*Monterey pine *P. radiata* Pm*red pine *P. resinosa* Pr*sugar pine *P. lambertiana* Ps

Spruce Picea S

*Norway spruce *P. abies* Sn

Changes to Version 4 of B.C. Ministry of Forests Tree Code List

1. Exotic species added to list: a) to provide codes for database purposes (requested by Resources Inventory Branch); and b) to accommodate inventories being conducted near areas of settlement that may encounter escaped or naturalized exotics.

2. Hybrids that cannot be easily distinguished or are of doubtful existence were deleted from list. Operational option is to go to upper level generic code if suspected hybrids without codes are encountered; common hybrids still have codes. Deletions include: Sxe (*Picea engelmannii* x *sitchensis*), Sxb (*Picea glauca* x *mariana*), Sxx (*Picea glauca* x *engelmannii* x *sitchensis*), and Exw (*Betula occidentalis* x *papyrifera*).

3. Dm (*Alnus tenuifolia*) deleted as it is not known to exceed 10 m in height and is most often multi-stemmed.

Changes to Version 4.1 of B.C. Ministry of Forests Tree Code List

Sn (*Picea abies*) Norway Spruce was added. Note that some printed copies of version 4.0 had Norway spruce included but our master version required the update hence the minor version upgrade.

Changes to Version 4.2 of B.C. Ministry of Forests Tree Code List

Four exotic species requiring codes for database purposes were added: yellow birch (Ey), white ash (Oh), shagbark hickory (Oi), and white oak (Qw).

Changes to Version 4.3 of B.C. Ministry of Forests Tree Code List

One exotic species requiring a code for database purposes was added: Dahurian larch (Ld).

Species Composition Code - Second Species

Input Format: XXX
Input Example: PL
Data Origin: input
Attribute Source: both

Sequence:	
Optional:	
Format:	varchar2
Length:	4
Decimal Places:	
Null:	Y

Use: The species code is used in determining: species composition, stand volumes, stand decay, waste and breakage, net-downs in Timber Supply Analyses, site index, etc.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_cd_3 Short Name: spec_cd_3
 Alias: species cd 3

Forestry Term: Species Composition Code - Third Species

Description: A code describing the leading commercial species or brush species in the layer. The species with the highest percent composition (e.g. gross volume or, if a very young stand, the relative number of stems per hectare) is identified as the leading commercial species. Species must be above a specified diameter to be recognized in the species composition of the layer. Leading species are described in terms of Genus, Species and Subspecies. There are currently 27 commercial tree species and five genus values recognized in the Province. The code may also be used to describe brush species in cases where the Non-Productive Descriptor is NPBR or the Non-Forest Descriptor is NCBR.

Measurement Criteria

Standard: 3 character alpha code indicating commercial species.

Default: blank

Permitted Values <blank> No species recorded

Commercial Species

AC	Balsam poplar	Populus balsamifera Black
	Cottonwood	Populus balsamifera
AT	Aspen	Populus tremuloides
B	True fir	Abies spp.
BL	Alpine fir	Abies lasiocarpa
BA	Amabilis fir	Abies amabilis
BG	Grand fir	Abies grandis
CW	Western red cedar	Thuja plicata
DR	Red Alder	Alnus rubra
E	Birch	Betula spp.
EP	Common paper birch	Betula papyrifera
EA	Alaska paper birch	Betula neoalaskana
FD	Douglas fir	Pseudotsuga menziesii
H	Hemlocks	Tsuga spp.
HW	Western hemlock	Tsuga heterophylla
HM	Mountain hemlock	Tsuga mertensiana
L	Larch	Larix spp.
LA	Alpine larch	Larix lyalli
LT	Tamarack	Larix laricina
LW	Western larch	Larix occidentalis
MB	Broadleaf maple	Acer macrophyllum
PF	Limber pine	Pinus flexilis
PL	Lodgepole pine	Pinus contorta
PW	Western white pine	Pinus monticola
PA	Whitebark pine	Pinus albicalis
PY	Yellow pine	Pinus ponderosa
PJ	Jack pine	Pinus banksiana
S	Spruce	Picea spp.
SB	Black spruce	Picea mariana
SE	Engelmann spruce	Picea engelmannii
SS	Sitka spruce	Picea sitchensis
SW	White spruce	Picea glauca
YC	Yellow cedar	Chamaecyparis nootkatensis

Brush Species

DM	Mountain alder	Alnus incana
R	Arbutus	Arbutus menziesii
EW	Water birch	Betula occidentalis

Cedar Thuja C

western redcedar Thuja plicata Cw

Cypress Chamaecyparis Y

yellow-cedar C. nootkatensis Yc

Douglas-fir *Pseudotsuga* F
 Douglas-fir *P. menziesii* Fd
 coastal Douglas-fir *P. menziesii* var. *menziesii* Fdc
 interior Douglas-fir *P. menziesii* var. *glauca* Fdi

Fir (Balsam) *Abies* B
 amabilis fir *A. amabilis* Ba
 grand fir *A. grandis* Bg
 subalpine fir *A. lasiocarpa* BI

Hemlock *Tsuga* H
 mountain hemlock *T. mertensiana* Hm
 western hemlock *T. heterophylla* Hw
 mountain x western hemlock hybrid *T. mertensiana* x *heterophylla* Hxm

Juniper *Juniperus* J
 Rocky Mtn. juniper *J. scopulorum* Jr

Larch *Larix* L
 alpine larch *L. lyallii* La
 tamarack *L. laricina* Lt
 western larch *L. occidentalis* Lw

Pine *Pinus* P
 jack pine *P. banksiana* Pj
 limber pine *P. flexilis* Pf
 lodgepole pine *P. contorta* Pl
 lodgepole pine *P. contorta* var. *latifolia* Pli
 lodgepole x jack pine hybrid *P. x murraybanksiana* Pxi
 ponderosa pine *P. ponderosa* Py
 shore pine *P. contorta* var. *contorta* Plc
 western white pine *P. monticola* Pw
 whitebark pine *P. albicaulis* Pa

Spruce *Picea* S
 black spruce *P. mariana* Sb
 Engelmann spruce *P. engelmannii* Se
 Sitka spruce *P. sitchensis* Ss
 white spruce *P. glauca* Sw
 spruce hybrid *Picea* cross Sx
 Engelmann x white *P. engelmannii* x *glauca* Sxw
 Sitka x white *P. x lutzii* Sxl
 Sitka x unknown hybrid *P. sitchensis* x ? Sxs

Yew *Taxus* T
 western yew *Taxus brevifolia* Tw

NATIVE HARDWOODS

Alder *Alnus* D
 red alder *A. rubra* Dr

Apple *Malus* U
 Pacific crab apple *Malus fusca* Up

Aspen, Cottonwood or Poplar *Populus* A
 poplar *P. balsamifera* Ac
 balsam poplar *P. b. ssp. balsamifera* Acb
 black cottonwood *P. b. ssp. trichocarpa* Act
 hybrid poplars *P. spp.* Ax
 trembling aspen *P. tremuloides* At

Arbutus *Arbutus* R
 Arbutus *Arbutus menziesii* Ra

Birch *Betula* E
 Alaska paper birch *B. neoalaskana* Ea
 Alaska x paper birch hybrid *B. x winteri* Exp

paper birch *B. papyrifera* Ep
water birch *B. occidentalis* Ew

Cascara *Rhamnus* K
cascara *R. purshiana* Kc

Cherry *Prunus* V
bitter cherry *P. emarginata* Vb
choke cherry *P. virginiana* Vv
pin cherry *P. pensylvanica* Vp

Dogwood *Cornus* G
Pacific dogwood *Cornus nuttallii* Gp

Maple *Acer* M
bigleaf maple *A. macrophyllum* Mb
vine maple *A. circinatum* Mv

Oak *Quercus* Q
Garry oak *Q. garryana* Qg

Willow *Salix* spp. W
Bebb's willow *S. bebbiana* Wb
Pacific willow *S. lucida* Wp
peachleaf willow *S. amygdaloides* Wa
pussy willow *S. discolor* Wd
Scouler's willow *S. scouleriana* Ws
Sitka willow *S. sitchensis* Wt

UNKNOWNNS

Unknown X
Unknown conifer Xc
Unknown hardwood Xh

OTHERS

Other tree, not on list Z
Other conifer Zc
Other hardwood Zh

EXOTICS

Apple *Malus* U
apple *Malus pumila* Ua

Aspen, Cottonwood or Poplar *Populus* A
*southern cottonwood *P. deltoides* Ad

Birch *Betula* E
European birch *B. pendula* Ee
silver birch *B. pubescens* Es
*yellow birch *B. alleghaniensis* Ey

Cherry *Prunus* V
sweet cherry *P. avium* Vs

Cypress *Chamaecyparis* Y
*Port Orford-cedar *C. lawsoniana* Yp

Fir (Balsam) *Abies* B
*balsam fir *A. balsamea* Bb
noble fir *A. procera* Bp
*Shasta red fir *A. magnifica* var. *shastensis* Bm
*white fir *A. concolor* Bc

Larch *Larix* L

*Dahurian larch *L. gmelinii* Ld

Maple Acer M

box elder *A. negundo* Me

*Norway maple *A. platanoides* Mn

*Sycamore maple *A. pseudoplatanus* Ms

Oak Quercus Q

*English oak *Q. robur* Qe

*white oak *Q. alba* Qw

Other exotics

*incense-cedar *Calocedrus decurrens* Oa

*giant sequoia *Sequoiadendron giganteum* Ob

*coast redwood *Sequoia sempervirens* Oc

European mountain-ash *Sorbus aucuparia* Od

Siberian elm *Ulmus pumila* Oe

common pear *Pyrus communis* Of

Oregon ash *Fraxinus latifolia* Og

*white ash *Fraxinus americana* Oh

*shagbark hickory *Carya ovata* Oi

Pine Pinus P

*Monterey pine *P. radiata* Pm

*red pine *P. resinosa* Pr

*sugar pine *P. lambertiana* Ps

Spruce Picea S

*Norway spruce *P. abies* Sn

Changes to Version 4 of B.C. Ministry of Forests Tree Code List

1. Exotic species added to list: a) to provide codes for database purposes (requested by Resources Inventory Branch); and b) to accommodate inventories being conducted near areas of settlement that may encounter escaped or naturalized exotics.

2. Hybrids that cannot be easily distinguished or are of doubtful existence were deleted from list. Operational option is to go to upper level generic code if suspected hybrids without codes are encountered; common hybrids still have codes. Deletions include: Sxe (*Picea engelmannii* x *sitchensis*), Sxb (*Picea glauca* x *mariana*), Sxx (*Picea glauca* x *engelmannii* x *sitchensis*), and Exw (*Betula occidentalis* x *papyrifera*).

3. Dm (*Alnus tenuifolia*) deleted as it is not known to exceed 10 m in height and is most often multi-stemmed.

Changes to Version 4.1 of B.C. Ministry of Forests Tree Code List

Sn (*Picea abies*) Norway Spruce was added. Note that some printed copies of version 4.0 had Norway spruce included but our master version required the update hence the minor version upgrade.

Changes to Version 4.2 of B.C. Ministry of Forests Tree Code List

Four exotic species requiring codes for database purposes were added: yellow birch (Ey), white ash (Oh), shagbark hickory (Oi), and white oak (Qw).

Changes to Version 4.3 of B.C. Ministry of Forests Tree Code List

One exotic species requiring a code for database purposes was added: Dahurian larch (Ld).

Species Composition Code - Third Species

Input Format: XXX
Input Example: PL
Data Origin: input
Attribute Source: both

Sequence:	
Optional:	
Format:	varchar2
Length:	4
Decimal Places:	
Null:	Y

Use: The species code is used in determining: species composition, stand volumes, stand decay, waste and breakage, net-downs in Timber Supply Analyses, site index, etc.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_cd_4 Short Name: spec_cd_4
 Alias: species cd 4

Forestry Term: Species Composition Code - Fourth Species

Description: A code describing the leading commercial species or brush species in the layer. The species with the highest percent composition (e.g. gross volume or, if a very young stand, the relative number of stems per hectare) is identified as the leading commercial species. Species must be above a specified diameter to be recognized in the species composition of the layer. Leading species are described in terms of Genus, Species and Subspecies. There are currently 27 commercial tree species and five genus values recognized in the Province. The code may also be used to describe brush species in cases where the Non-Productive Descriptor is NPBR or the Non-Forest Descriptor is NCBR.

Measurement Criteria

Standard: 3 character alpha code indicating commercial species.

Default: blank

Permitted Values <blank> No species recorded

Commercial Species

AC	Balsam poplar	Populus balsamifera Black
	Cottonwood	Populus balsamifera
AT	Aspen	Populus tremuloides
B	True fir	Abies spp.
BL	Alpine fir	Abies lasiocarpa
BA	Amabilis fir	Abies amabilis
BG	Grand fir	Abies grandis
CW	Western red cedar	Thuja plicata
DR	Red Alder	Alnus rubra
E	Birch	Betula spp.
EP	Common paper birch	Betula papyrifera
EA	Alaska paper birch	Betula neoalaskana
FD	Douglas fir	Pseudotsuga menziesii
H	Hemlocks	Tsuga spp.
HW	Western hemlock	Tsuga heterophylla
HM	Mountain hemlock	Tsuga mertensiana
L	Larch	Larix spp.
LA	Alpine larch	Larix lyalli
LT	Tamarack	Larix laricina
LW	Western larch	Larix occidentalis
MB	Broadleaf maple	Acer macrophyllum
PF	Limber pine	Pinus flexilis
PL	Lodgepole pine	Pinus contorta
PW	Western white pine	Pinus monticola
PA	Whitebark pine	Pinus albicalis
PY	Yellow pine	Pinus ponderosa
PJ	Jack pine	Pinus banksiana
S	Spruce	Picea spp.
SB	Black spruce	Picea mariana
SE	Engelmann spruce	Picea engelmannii
SS	Sitka spruce	Picea sitchensis
SW	White spruce	Picea glauca
YC	Yellow cedar	Chamaecyparis nootkatensis

Brush Species

DM	Mountain alder	Alnus incana
R	Arbutus	Arbutus menziesii
EW	Water birch	Betula occidentalis

Cedar Thuja C

western redcedar Thuja plicata Cw

Cypress Chamaecyparis Y

yellow-cedar C. nootkatensis Yc

Douglas-fir *Pseudotsuga* F
 Douglas-fir *P. menziesii* Fd
 coastal Douglas-fir *P. menziesii* var. *menziesii* Fdc
 interior Douglas-fir *P. menziesii* var. *glauca* Fdi

Fir (Balsam) *Abies* B
 amabilis fir *A. amabilis* Ba
 grand fir *A. grandis* Bg
 subalpine fir *A. lasiocarpa* BI

Hemlock *Tsuga* H
 mountain hemlock *T. mertensiana* Hm
 western hemlock *T. heterophylla* Hw
 mountain x western hemlock hybrid *T. mertensiana* x *heterophylla* Hxm

Juniper *Juniperus* J
 Rocky Mtn. juniper *J. scopulorum* Jr

Larch *Larix* L
 alpine larch *L. lyallii* La
 tamarack *L. laricina* Lt
 western larch *L. occidentalis* Lw

Pine *Pinus* P
 jack pine *P. banksiana* Pj
 limber pine *P. flexilis* Pf
 lodgepole pine *P. contorta* Pl
 lodgepole pine *P. contorta* var. *latifolia* Pli
 lodgepole x jack pine hybrid *P. x murraybanksiana* Pxi
 ponderosa pine *P. ponderosa* Py
 shore pine *P. contorta* var. *contorta* Plc
 western white pine *P. monticola* Pw
 whitebark pine *P. albicaulis* Pa

Spruce *Picea* S
 black spruce *P. mariana* Sb
 Engelmann spruce *P. engelmannii* Se
 Sitka spruce *P. sitchensis* Ss
 white spruce *P. glauca* Sw
 spruce hybrid *Picea* cross Sx
 Engelmann x white *P. engelmannii* x *glauca* Sxw
 Sitka x white *P. x lutzii* Sxl
 Sitka x unknown hybrid *P. sitchensis* x ? Sxs

Yew *Taxus* T
 western yew *Taxus brevifolia* Tw

NATIVE HARDWOODS

Alder *Alnus* D
 red alder *A. rubra* Dr

Apple *Malus* U
 Pacific crab apple *Malus fusca* Up

Aspen, Cottonwood or Poplar *Populus* A
 poplar *P. balsamifera* Ac
 balsam poplar *P. b. ssp. balsamifera* Acb
 black cottonwood *P. b. ssp. trichocarpa* Act
 hybrid poplars *P. spp.* Ax
 trembling aspen *P. tremuloides* At

Arbutus *Arbutus* R
 Arbutus *Arbutus menziesii* Ra

Birch *Betula* E
 Alaska paper birch *B. neoalaskana* Ea
 Alaska x paper birch hybrid *B. x winteri* Exp

paper birch *B. papyrifera* Ep
water birch *B. occidentalis* Ew

Cascara *Rhamnus* K
cascara *R. purshiana* Kc

Cherry *Prunus* V
bitter cherry *P. emarginata* Vb
choke cherry *P. virginiana* Vv
pin cherry *P. pensylvanica* Vp

Dogwood *Cornus* G
Pacific dogwood *Cornus nuttallii* Gp

Maple *Acer* M
bigleaf maple *A. macrophyllum* Mb
vine maple *A. circinatum* Mv

Oak *Quercus* Q
Garry oak *Q. garryana* Qg

Willow *Salix* spp. W
Bebb's willow *S. bebbiana* Wb
Pacific willow *S. lucida* Wp
peachleaf willow *S. amygdaloides* Wa
pussy willow *S. discolor* Wd
Scouler's willow *S. scouleriana* Ws
Sitka willow *S. sitchensis* Wt

UNKNOWNNS

Unknown X
Unknown conifer Xc
Unknown hardwood Xh

OTHERS

Other tree, not on list Z
Other conifer Zc
Other hardwood Zh

EXOTICS

Apple *Malus* U
apple *Malus pumila* Ua

Aspen, Cottonwood or Poplar *Populus* A
*southern cottonwood *P. deltoides* Ad

Birch *Betula* E
European birch *B. pendula* Ee
silver birch *B. pubescens* Es
*yellow birch *B. alleghaniensis* Ey

Cherry *Prunus* V
sweet cherry *P. avium* Vs

Cypress *Chamaecyparis* Y
*Port Orford-cedar *C. lawsoniana* Yp

Fir (Balsam) *Abies* B
*balsam fir *A. balsamea* Bb
noble fir *A. procera* Bp
*Shasta red fir *A. magnifica* var. *shastensis* Bm
*white fir *A. concolor* Bc

Larch *Larix* L

*Dahurian larch *L. gmelinii* Ld

Maple Acer M

box elder *A. negundo* Me*Norway maple *A. platanoides* Mn*Sycamore maple *A. pseudoplatanus* Ms

Oak Quercus Q

*English oak *Q. robur* Qe*white oak *Q. alba* Qw

Other exotics

*incense-cedar *Calocedrus decurrens* Oa*giant sequoia *Sequoiadendron giganteum* Ob*coast redwood *Sequoia sempervirens* OcEuropean mountain-ash *Sorbus aucuparia* OdSiberian elm *Ulmus pumila* Oecommon pear *Pyrus communis* OfOregon ash *Fraxinus latifolia* Og*white ash *Fraxinus americana* Oh*shagbark hickory *Carya ovata* Oi

Pine Pinus P

*Monterey pine *P. radiata* Pm*red pine *P. resinosa* Pr*sugar pine *P. lambertiana* Ps

Spruce Picea S

*Norway spruce *P. abies* Sn

Changes to Version 4 of B.C. Ministry of Forests Tree Code List

1. Exotic species added to list: a) to provide codes for database purposes (requested by Resources Inventory Branch); and b) to accommodate inventories being conducted near areas of settlement that may encounter escaped or naturalized exotics.

2. Hybrids that cannot be easily distinguished or are of doubtful existence were deleted from list. Operational option is to go to upper level generic code if suspected hybrids without codes are encountered; common hybrids still have codes. Deletions include: Sxe (*Picea engelmannii* x *sitchensis*), Sxb (*Picea glauca* x *mariana*), Sxx (*Picea glauca* x *engelmannii* x *sitchensis*), and Exw (*Betula occidentalis* x *papyrifera*).

3. Dm (*Alnus tenuifolia*) deleted as it is not known to exceed 10 m in height and is most often multi-stemmed.

Changes to Version 4.1 of B.C. Ministry of Forests Tree Code List

Sn (*Picea abies*) Norway Spruce was added. Note that some printed copies of version 4.0 had Norway spruce included but our master version required the update hence the minor version upgrade.

Changes to Version 4.2 of B.C. Ministry of Forests Tree Code List

Four exotic species requiring codes for database purposes were added: yellow birch (Ey), white ash (Oh), shagbark hickory (Oi), and white oak (Qw).

Changes to Version 4.3 of B.C. Ministry of Forests Tree Code List

One exotic species requiring a code for database purposes was added: Dahurian larch (Ld).

Species Composition Code - Fourth Species

Input Format: XXX
Input Example: PL
Data Origin: input
Attribute Source: both

Sequence:	
Optional:	
Format:	varchar2
Length:	4
Decimal Places:	
Null:	Y

Use: The species code is used in determining: species composition, stand volumes, stand decay, waste and breakage, net-downs in Timber Supply Analyses, site index, etc.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_cd_5 Short Name: spec_cd_5
 Alias: species cd 5

Forestry Term: Species Composition Code - Fifth Species

Description: A code describing the leading commercial species or brush species in the layer. The species with the highest percent composition (e.g. gross volume or, if a very young stand, the relative number of stems per hectare) is identified as the leading commercial species. Species must be above a specified diameter to be recognized in the species composition of the layer. Leading species are described in terms of Genus, Species and Subspecies. There are currently 27 commercial tree species and five genus values recognized in the Province. The code may also be used to describe brush species in cases where the Non-Productive Descriptor is NPBR or the Non-Forest Descriptor is NCBR.

Measurement Criteria

Standard: 3 character alpha code indicating commercial species.

Default: blank

Permitted Values <blank> No species recorded

Commercial Species

AC	Balsam poplar	Populus balsamifera Black
	Cottonwood	Populus balsamifera
AT	Aspen	Populus tremuloides
B	True fir	Abies spp.
BL	Alpine fir	Abies lasiocarpa
BA	Amabilis fir	Abies amabilis
BG	Grand fir	Abies grandis
CW	Western red cedar	Thuja plicata
DR	Red Alder	Alnus rubra
E	Birch	Betula spp.
EP	Common paper birch	Betula papyrifera
EA	Alaska paper birch	Betula neoalaskana
FD	Douglas fir	Pseudotsuga menziesii
H	Hemlocks	Tsuga spp.
HW	Western hemlock	Tsuga heterophylla
HM	Mountain hemlock	Tsuga mertensiana
L	Larch	Larix spp.
LA	Alpine larch	Larix lyalli
LT	Tamarack	Larix laricina
LW	Western larch	Larix occidentalis
MB	Broadleaf maple	Acer macrophyllum
PF	Limber pine	Pinus flexilis
PL	Lodgepole pine	Pinus contorta
PW	Western white pine	Pinus monticola
PA	Whitebark pine	Pinus albicalis
PY	Yellow pine	Pinus ponderosa
PJ	Jack pine	Pinus banksiana
S	Spruce	Picea spp.
SB	Black spruce	Picea mariana
SE	Engelmann spruce	Picea engelmannii
SS	Sitka spruce	Picea sitchensis
SW	White spruce	Picea glauca
YC	Yellow cedar	Chamaecyparis nootkatensis

Brush Species

DM	Mountain alder	Alnus incana
R	Arbutus	Arbutus menziesii
EW	Water birch	Betula occidentalis

Cedar Thuja C

western redcedar Thuja plicata Cw

Cypress Chamaecyparis Y

yellow-cedar C. nootkatensis Yc

Douglas-fir *Pseudotsuga* F
 Douglas-fir *P. menziesii* Fd
 coastal Douglas-fir *P. menziesii* var. *menziesii* Fdc
 interior Douglas-fir *P. menziesii* var. *glauca* Fdi

Fir (Balsam) *Abies* B
 amabilis fir *A. amabilis* Ba
 grand fir *A. grandis* Bg
 subalpine fir *A. lasiocarpa* BI

Hemlock *Tsuga* H
 mountain hemlock *T. mertensiana* Hm
 western hemlock *T. heterophylla* Hw
 mountain x western hemlock hybrid *T. mertensiana* x *heterophylla* Hxm

Juniper *Juniperus* J
 Rocky Mtn. juniper *J. scopulorum* Jr

Larch *Larix* L
 alpine larch *L. lyallii* La
 tamarack *L. laricina* Lt
 western larch *L. occidentalis* Lw

Pine *Pinus* P
 jack pine *P. banksiana* Pj
 limber pine *P. flexilis* Pf
 lodgepole pine *P. contorta* Pl
 lodgepole pine *P. contorta* var. *latifolia* Pli
 lodgepole x jack pine hybrid *P. x murraybanksiana* Pxi
 ponderosa pine *P. ponderosa* Py
 shore pine *P. contorta* var. *contorta* Plc
 western white pine *P. monticola* Pw
 whitebark pine *P. albicaulis* Pa

Spruce *Picea* S
 black spruce *P. mariana* Sb
 Engelmann spruce *P. engelmannii* Se
 Sitka spruce *P. sitchensis* Ss
 white spruce *P. glauca* Sw
 spruce hybrid *Picea* cross Sx
 Engelmann x white *P. engelmannii* x *glauca* Sxw
 Sitka x white *P. x lutzii* Sxl
 Sitka x unknown hybrid *P. sitchensis* x ? Sxs

Yew *Taxus* T
 western yew *Taxus brevifolia* Tw

NATIVE HARDWOODS

Alder *Alnus* D
 red alder *A. rubra* Dr

Apple *Malus* U
 Pacific crab apple *Malus fusca* Up

Aspen, Cottonwood or Poplar *Populus* A
 poplar *P. balsamifera* Ac
 balsam poplar *P. b. ssp. balsamifera* Acb
 black cottonwood *P. b. ssp. trichocarpa* Act
 hybrid poplars *P. spp.* Ax
 trembling aspen *P. tremuloides* At

Arbutus *Arbutus* R
 Arbutus *Arbutus menziesii* Ra

Birch *Betula* E
 Alaska paper birch *B. neoalaskana* Ea
 Alaska x paper birch hybrid *B. x winteri* Exp

paper birch *B. papyrifera* Ep
water birch *B. occidentalis* Ew

Cascara *Rhamnus* K
cascara *R. purshiana* Kc

Cherry *Prunus* V
bitter cherry *P. emarginata* Vb
choke cherry *P. virginiana* Vv
pin cherry *P. pensylvanica* Vp

Dogwood *Cornus* G
Pacific dogwood *Cornus nuttallii* Gp

Maple *Acer* M
bigleaf maple *A. macrophyllum* Mb
vine maple *A. circinatum* Mv

Oak *Quercus* Q
Garry oak *Q. garryana* Qg

Willow *Salix* spp. W
Bebb's willow *S. bebbiana* Wb
Pacific willow *S. lucida* Wp
peachleaf willow *S. amygdaloides* Wa
pussy willow *S. discolor* Wd
Scouler's willow *S. scouleriana* Ws
Sitka willow *S. sitchensis* Wt

UNKNOWNNS

Unknown X
Unknown conifer Xc
Unknown hardwood Xh

OTHERS

Other tree, not on list Z
Other conifer Zc
Other hardwood Zh

EXOTICS

Apple *Malus* U
apple *Malus pumila* Ua

Aspen, Cottonwood or Poplar *Populus* A
*southern cottonwood *P. deltoides* Ad

Birch *Betula* E
European birch *B. pendula* Ee
silver birch *B. pubescens* Es
*yellow birch *B. alleghaniensis* Ey

Cherry *Prunus* V
sweet cherry *P. avium* Vs

Cypress *Chamaecyparis* Y
*Port Orford-cedar *C. lawsoniana* Yp

Fir (Balsam) *Abies* B
*balsam fir *A. balsamea* Bb
noble fir *A. procera* Bp
*Shasta red fir *A. magnifica* var. *shastensis* Bm
*white fir *A. concolor* Bc

Larch *Larix* L

*Dahurian larch *L. gmelinii* Ld

Maple Acer M

box elder *A. negundo* Me*Norway maple *A. platanoides* Mn*Sycamore maple *A. pseudoplatanus* Ms

Oak Quercus Q

*English oak *Q. robur* Qe*white oak *Q. alba* Qw

Other exotics

*incense-cedar *Calocedrus decurrens* Oa*giant sequoia *Sequoiadendron giganteum* Ob*coast redwood *Sequoia sempervirens* OcEuropean mountain-ash *Sorbus aucuparia* OdSiberian elm *Ulmus pumila* Oecommon pear *Pyrus communis* OfOregon ash *Fraxinus latifolia* Og*white ash *Fraxinus americana* Oh*shagbark hickory *Carya ovata* Oi

Pine Pinus P

*Monterey pine *P. radiata* Pm*red pine *P. resinosa* Pr*sugar pine *P. lambertiana* Ps

Spruce Picea S

*Norway spruce *P. abies* Sn

Changes to Version 4 of B.C. Ministry of Forests Tree Code List

1. Exotic species added to list: a) to provide codes for database purposes (requested by Resources Inventory Branch); and b) to accommodate inventories being conducted near areas of settlement that may encounter escaped or naturalized exotics.

2. Hybrids that cannot be easily distinguished or are of doubtful existence were deleted from list. Operational option is to go to upper level generic code if suspected hybrids without codes are encountered; common hybrids still have codes. Deletions include: Sxe (*Picea engelmannii* x *sitchensis*), Sxb (*Picea glauca* x *mariana*), Sxx (*Picea glauca* x *engelmannii* x *sitchensis*), and Exw (*Betula occidentalis* x *papyrifera*).

3. Dm (*Alnus tenuifolia*) deleted as it is not known to exceed 10 m in height and is most often multi-stemmed.

Changes to Version 4.1 of B.C. Ministry of Forests Tree Code List

Sn (*Picea abies*) Norway Spruce was added. Note that some printed copies of version 4.0 had Norway spruce included but our master version required the update hence the minor version upgrade.

Changes to Version 4.2 of B.C. Ministry of Forests Tree Code List

Four exotic species requiring codes for database purposes were added: yellow birch (Ey), white ash (Oh), shagbark hickory (Oi), and white oak (Qw).

Changes to Version 4.3 of B.C. Ministry of Forests Tree Code List

One exotic species requiring a code for database purposes was added: Dahurian larch (Ld).

Input Format: XXX
Input Example: PL
Data Origin: input
Attribute Source: both

Sequence:	
Optional:	
Format:	varchar2
Length:	4
Decimal Places:	
Null:	Y

Use: The species code is used in determining: species composition, stand volumes, stand decay, waste and breakage, net-downs in Timber Supply Analyses, site index, etc.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_cd_6 Short Name: spec_cd_6
 Alias: species cd 6

Forestry Term: Species Composition Code - Sixth Species

Description: A code describing the leading commercial species or brush species in the layer. The species with the highest percent composition (e.g. gross volume or, if a very young stand, the relative number of stems per hectare) is identified as the leading commercial species. Species must be above a specified diameter to be recognized in the species composition of the layer. Leading species are described in terms of Genus, Species and Subspecies. There are currently 27 commercial tree species and five genus values recognized in the Province. The code may also be used to describe brush species in cases where the Non-Productive Descriptor is NPBR or the Non-Forest Descriptor is NCBR.

Measurement Criteria

Standard: 3 character alpha code indicating commercial species.

Default: blank

Permitted Values <blank> No species recorded

Commercial Species

AC	Balsam poplar	Populus balsamifera Black
	Cottonwood	Populus balsamifera
AT	Aspen	Populus tremuloides
B	True fir	Abies spp.
BL	Alpine fir	Abies lasiocarpa
BA	Amabilis fir	Abies amabilis
BG	Grand fir	Abies grandis
CW	Western red cedar	Thuja plicata
DR	Red Alder	Alnus rubra
E	Birch	Betula spp.
EP	Common paper birch	Betula papyrifera
EA	Alaska paper birch	Betula neoalaskana
FD	Douglas fir	Pseudotsuga menziesii
H	Hemlocks	Tsuga spp.
HW	Western hemlock	Tsuga heterophylla
HM	Mountain hemlock	Tsuga mertensiana
L	Larch	Larix spp.
LA	Alpine larch	Larix lyalli
LT	Tamarack	Larix laricina
LW	Western larch	Larix occidentalis
MB	Broadleaf maple	Acer macrophyllum
PF	Limber pine	Pinus flexilis
PL	Lodgepole pine	Pinus contorta
PW	Western white pine	Pinus monticola
PA	Whitebark pine	Pinus albicalis
PY	Yellow pine	Pinus ponderosa
PJ	Jack pine	Pinus banksiana
S	Spruce	Picea spp.
SB	Black spruce	Picea mariana
SE	Engelmann spruce	Picea engelmannii
SS	Sitka spruce	Picea sitchensis
SW	White spruce	Picea glauca
YC	Yellow cedar	Chamaecyparis nootkatensis

Brush Species

DM	Mountain alder	Alnus incana
R	Arbutus	Arbutus menziesii
EW	Water birch	Betula occidentalis

Cedar Thuja C

western redcedar Thuja plicata Cw

Cypress Chamaecyparis Y

yellow-cedar C. nootkatensis Yc

Douglas-fir *Pseudotsuga* F
 Douglas-fir *P. menziesii* Fd
 coastal Douglas-fir *P. menziesii* var. *menziesii* Fdc
 interior Douglas-fir *P. menziesii* var. *glauca* Fdi

Fir (Balsam) *Abies* B
 amabilis fir *A. amabilis* Ba
 grand fir *A. grandis* Bg
 subalpine fir *A. lasiocarpa* BI

Hemlock *Tsuga* H
 mountain hemlock *T. mertensiana* Hm
 western hemlock *T. heterophylla* Hw
 mountain x western hemlock hybrid *T. mertensiana* x *heterophylla* Hxm

Juniper *Juniperus* J
 Rocky Mtn. juniper *J. scopulorum* Jr

Larch *Larix* L
 alpine larch *L. lyallii* La
 tamarack *L. laricina* Lt
 western larch *L. occidentalis* Lw

Pine *Pinus* P
 jack pine *P. banksiana* Pj
 limber pine *P. flexilis* Pf
 lodgepole pine *P. contorta* Pl
 lodgepole pine *P. contorta* var. *latifolia* Pli
 lodgepole x jack pine hybrid *P. x murraybanksiana* Pxi
 ponderosa pine *P. ponderosa* Py
 shore pine *P. contorta* var. *contorta* Plc
 western white pine *P. monticola* Pw
 whitebark pine *P. albicaulis* Pa

Spruce *Picea* S
 black spruce *P. mariana* Sb
 Engelmann spruce *P. engelmannii* Se
 Sitka spruce *P. sitchensis* Ss
 white spruce *P. glauca* Sw
 spruce hybrid *Picea* cross Sx
 Engelmann x white *P. engelmannii* x *glauca* Sxw
 Sitka x white *P. x lutzii* Sxl
 Sitka x unknown hybrid *P. sitchensis* x ? Sxs

Yew *Taxus* T
 western yew *Taxus brevifolia* Tw

NATIVE HARDWOODS

Alder *Alnus* D
 red alder *A. rubra* Dr

Apple *Malus* U
 Pacific crab apple *Malus fusca* Up

Aspen, Cottonwood or Poplar *Populus* A
 poplar *P. balsamifera* Ac
 balsam poplar *P. b. ssp. balsamifera* Acb
 black cottonwood *P. b. ssp. trichocarpa* Act
 hybrid poplars *P. spp.* Ax
 trembling aspen *P. tremuloides* At

Arbutus *Arbutus* R
 Arbutus *Arbutus menziesii* Ra

Birch *Betula* E
 Alaska paper birch *B. neoalaskana* Ea
 Alaska x paper birch hybrid *B. x winteri* Exp

paper birch *B. papyrifera* Ep
water birch *B. occidentalis* Ew

Cascara *Rhamnus* K
cascara *R. purshiana* Kc

Cherry *Prunus* V
bitter cherry *P. emarginata* Vb
choke cherry *P. virginiana* Vv
pin cherry *P. pensylvanica* Vp

Dogwood *Cornus* G
Pacific dogwood *Cornus nuttallii* Gp

Maple *Acer* M
bigleaf maple *A. macrophyllum* Mb
vine maple *A. circinatum* Mv

Oak *Quercus* Q
Garry oak *Q. garryana* Qg

Willow *Salix* spp. W
Bebb's willow *S. bebbiana* Wb
Pacific willow *S. lucida* Wp
peachleaf willow *S. amygdaloides* Wa
pussy willow *S. discolor* Wd
Scouler's willow *S. scouleriana* Ws
Sitka willow *S. sitchensis* Wt

UNKNOWNNS

Unknown X
Unknown conifer Xc
Unknown hardwood Xh

OTHERS

Other tree, not on list Z
Other conifer Zc
Other hardwood Zh

EXOTICS

Apple *Malus* U
apple *Malus pumila* Ua

Aspen, Cottonwood or Poplar *Populus* A
*southern cottonwood *P. deltoides* Ad

Birch *Betula* E
European birch *B. pendula* Ee
silver birch *B. pubescens* Es
*yellow birch *B. alleghaniensis* Ey

Cherry *Prunus* V
sweet cherry *P. avium* Vs

Cypress *Chamaecyparis* Y
*Port Orford-cedar *C. lawsoniana* Yp

Fir (Balsam) *Abies* B
*balsam fir *A. balsamea* Bb
noble fir *A. procera* Bp
*Shasta red fir *A. magnifica* var. *shastensis* Bm
*white fir *A. concolor* Bc

Larch *Larix* L

*Dahurian larch *L. gmelinii* Ld

Maple Acer M

box elder *A. negundo* Me*Norway maple *A. platanoides* Mn*Sycamore maple *A. pseudoplatanus* Ms

Oak Quercus Q

*English oak *Q. robur* Qe*white oak *Q. alba* Qw

Other exotics

*incense-cedar *Calocedrus decurrens* Oa*giant sequoia *Sequoiadendron giganteum* Ob*coast redwood *Sequoia sempervirens* OcEuropean mountain-ash *Sorbus aucuparia* OdSiberian elm *Ulmus pumila* Oecommon pear *Pyrus communis* OfOregon ash *Fraxinus latifolia* Og*white ash *Fraxinus americana* Oh*shagbark hickory *Carya ovata* Oi

Pine Pinus P

*Monterey pine *P. radiata* Pm*red pine *P. resinosa* Pr*sugar pine *P. lambertiana* Ps

Spruce Picea S

*Norway spruce *P. abies* Sn

Changes to Version 4 of B.C. Ministry of Forests Tree Code List

1. Exotic species added to list: a) to provide codes for database purposes (requested by Resources Inventory Branch); and b) to accommodate inventories being conducted near areas of settlement that may encounter escaped or naturalized exotics.

2. Hybrids that cannot be easily distinguished or are of doubtful existence were deleted from list. Operational option is to go to upper level generic code if suspected hybrids without codes are encountered; common hybrids still have codes. Deletions include: Sxe (*Picea engelmannii* x *sitchensis*), Sxb (*Picea glauca* x *mariana*), Sxx (*Picea glauca* x *engelmannii* x *sitchensis*), and Exw (*Betula occidentalis* x *papyrifera*).

3. Dm (*Alnus tenuifolia*) deleted as it is not known to exceed 10 m in height and is most often multi-stemmed.

Changes to Version 4.1 of B.C. Ministry of Forests Tree Code List

Sn (*Picea abies*) Norway Spruce was added. Note that some printed copies of version 4.0 had Norway spruce included but our master version required the update hence the minor version upgrade.

Changes to Version 4.2 of B.C. Ministry of Forests Tree Code List

Four exotic species requiring codes for database purposes were added: yellow birch (Ey), white ash (Oh), shagbark hickory (Oi), and white oak (Qw).

Changes to Version 4.3 of B.C. Ministry of Forests Tree Code List

One exotic species requiring a code for database purposes was added: Dahurian larch (Ld).

Input Format: XXX
Input Example: PL
Data Origin: input
Attribute Source: both

Sequence:	
Optional:	
Format:	varchar2
Length:	4
Decimal Places:	
Null:	Y

Use: The species code is used in determining: species composition, stand volumes, stand decay, waste and breakage, net-downs in Timber Supply Analyses, site index, etc.

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_pct_1 Short Name: spec_pct_1
Alias: species pct 1

Forestry Term: Leading Species Percentage

Description: Percentages of the layer that each tree species occupies. For older stands, tree species percentage is based on relative basal area; for younger stands, tree species percentage is based on the number of stems per hectare. Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Measurement Criteria: Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Standard: 3 character numeric value holding percent composition

Default: must have value, may be 0 if no species

Permitted Values: 0 to 100

Input Format: ###

Input Example: 60

Data Origin: input

Attribute Source: both

Sequence:	
Optional:	
Format:	number
Length:	3
Decimal Places:	
Null:	Y

Use: Tree species percentage is used in determining stand volumes, identifying stands with specific species composition (I.e. pure), net-downs in Timber Supply Analyses

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_pct_3 Short Name: spec_pct_3
Alias: species pct 3

Forestry Term: Third Species Percentage

Description: Percentages of the layer that each tree species occupies. For older stands, tree species percentage is based on relative basal area; for younger stands, tree species percentage is based on the number of stems per hectare. Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Measurement Criteria: Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Standard: 2 character numeric value holding percent composition

Default: must have value, may be 0 if no species

Permitted Values: 0 to 33

Input Format: ##
Input Example: 20
Data Origin: input
Attribute Source: both

Sequence:
Optional:
Format: number
Length: 2
Decimal Places:
Null: Y

Use: Tree species percentage is used in determining stand volumes, identifying stands with specific species composition (I.e. pure), net-downs in Timber Supply Analyses

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_pct_4 Short Name: spec_pct_4
 Alias: species pct 4

Forestry Term: Fourth Species Percentage

Description: Percentages of the layer that each tree species occupies. For older stands, tree species percentage is based on relative basal area; for younger stands, tree species percentage is based on the number of stems per hectare. Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Measurement Criteria: Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Standard: 2 character numeric value holding percent composition

Default: must have value, may be 0 if no species

Permitted Values: 0 to 25

Input Format: ##
 Input Example: 20
 Data Origin: input
 Attribute Source: both

Sequence:
 Optional:
 Format: number
 Length: 2
 Decimal Places:
 Null: Y

Use: Tree species percentage is used in determining stand volumes, identifying stands with specific species composition (I.e. pure), net-downs in Timber Supply Analyses

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_pct_5 Short Name: spec_pct_5
Alias: species pct 5

Forestry Term: Fifth Species Percentage

Description: Percentages of the layer that each tree species occupies. For older stands, tree species percentage is based on relative basal area; for younger stands, tree species percentage is based on the number of stems per hectare. Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Measurement Criteria: Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Standard: 2 character numeric value holding percent composition

Default: must have value, may be 0 if no species

Permitted Values: 0 to 20

Input Format: ##
Input Example: 10
Data Origin: input
Attribute Source: both

Sequence:
Optional:
Format: number
Length: 2
Decimal Places:
Null: Y

Use: Tree species percentage is used in determining stand volumes, identifying stands with specific species composition (I.e. pure), net-downs in Timber Supply Analyses

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: species_pct_6 Short Name: spec_pct_6
Alias: species pct 6

Forestry Term: Sixth Species Percentage

Description: Percentages of the layer that each tree species occupies. For older stands, tree species percentage is based on relative basal area; for younger stands, tree species percentage is based on the number of stems per hectare. Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Measurement Criteria: Tree species percentage is estimated to the nearest percent for all living trees above a specified diameter.

Standard: 2 character numeric value holding percent composition

Default: must have value, may be 0 if no species

Permitted Values: 0 to 16

Input Format: ##
Input Example: 10
Data Origin: input
Attribute Source: both

Sequence:
Optional:
Format: number
Length: 2
Decimal Places:
Null: Y

Use: Tree species percentage is used in determining stand volumes, identifying stands with specific species composition (I.e. pure), net-downs in Timber Supply Analyses

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: stand_percentage_dead Short Name: dead_pct
Alias stand percentage dead

Forestry Term: Stand Percentage Dead

Description: Represents the percent of the stand that has had an epidemic loss.

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###
Input Example: 70
Data Origin: input
Attribute Source:

Sequence:	
Optional:	
Format:	number
Length:	3
Decimal Places:	
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference:

Sub Type: veg_comp_poly

Attribute Name: surface_expression Short Name: surf_exp

Alias surface expression

Forestry Term: Surface Expression

Description: The form of surficial material apparent on the medium scale photography. A simplified attribution is used owing to the likelihood that the trees will mask surficial features.

Measurement Criteria Assign the appropriate letter code to each polygon.

Standard: In polygons that have multiple components, record the prevalent surface expression of the polygon on the basis of greatest percent area coverage.

Default:

Permitted Values Codes Description

C Cone

A cone, or segment of a cone, with a relatively smooth slope gradient, greater than 15 degrees (>25%).

D Depression

Circular or irregular area of lower elevation (such as a hollow) than the surrounding terrain; depressions are greater than two metres deep. Examples are kettle holes and karsts depressions

F Fan

A smooth segment of a cone with a slope gradient of up to 15 degrees (25%). Typically applied to fluvial or alluvial fans.

H Hummock(s)

Steep sided hillocks and hollows with slopes of 15 to 35 degrees (25 to 70%) predominant on unconsolidated materials, and slopes of 15 to 90 degrees (25% to vertical) predominant on bedrock. Slopes are non-linear (not parallel) but, generally, chaotic or dissected and rounded or irregular in profile. Local relief is greater than one metre. Differentiated from undulating on the basis of slope angle.

M Rolling

Elongated hillock(s) with slopes dominantly between 3 and 15 degrees (5 to 25%) with local relief greater than one metre. Slopes are an assemblage of parallel or sub-parallel linear forms with subdued relief and may occur in level or sloped meso slope positions.

N None of these descriptions apply as no apparent surface expression features are present.

P Plain

A level or gently sloping unidirectional surface with gradients of up to three degrees (5%). Local surface irregularities generally have a relief of less than one metre.

R Ridge(s)

Elongated or linear, parallel or sub-parallel hillock(s) or ridges with slopes predominantly between 15 and 35 degrees (25 to 70%) on unconsolidated materials and between 15 and 90 degrees (25% to vertical) on bedrock. Local relief is greater than one metre. Differentiated from rolling on the basis of slope angle. Possible locations include drumlinized till plains, eskers and ridged bedrock. These may be created through the erosional effects of water.

T Terrace(s)

Step-like topography where each step-like form consists of both a scarp face and a horizontal or gently inclined surface above it. The terrace description is applied to both the scarp and the flat surface.

U Undulating

Gently sloping hillock(s) and hollow(s) with slopes of up to 15 degrees (25%). Local relief is greater than one metre. Slopes are non-linear (not parallel), chaotic forms that are rounded or irregular in profile.

Input Format: X
 Input Example: N
 Data Origin: input
 Attribute Source: vri

Sequence:	30
Optional:	Y
Format:	varchar2
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_poly

Notes: Together with the attributes "modifying processes" and "site position meso" will provide clues to soil parent material and useful site classification data.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: tree_cover_pattern Short Name: tree_patrn

Alias: tree cover pattern

Forestry Term: Tree Cover Pattern

Description: A numeric code that identifies the spatial distribution of the tree layer in the polygon. Examples include treed islands in the sub-alpine parkland, clumps of trees on rocky outcrops, scattered groves or individual trees in an otherwise shrubby flood plain, or solid continuous tree cover. Tree cover pattern provides information on the amount of "edge" and "interior" habitat or growing conditions within the polygon. Tree cover pattern describes the spatial distribution of the tree cover within each tree layer in the polygon. Tree cover pattern is used to describe the tree layer spatial distribution.

Measurement Criteria: Cover pattern is estimated for each tree layer in the polygon.

Standard: Cover pattern is based on the majority area coverage.

Default:

- Permitted Values
1. Single to very few (<4) occurrences of limited extent, circular to irregular shape.
 2. Single to very few (<4) occurrences of limited extent, linear or elongated shape.
 3. Several (>3) sporadic occurrences of limited extent, circular to irregular shape.
 4. Several (>3) sporadic occurrences of limited extent, linear or elongated shape.
 5. Intimately intermixed units, often with gradational transitions from one to the other.
 6. Discontinuous but extensive occurrences, parallel to sub-parallel elongated in shape.
 7. Limited continuous occurrence with few inclusions.
 8. Continuous occurrence with several inclusions.
 9. Continuous occurrence with very few inclusions.

Input Format: #
 Input Example: 4
 Data Origin: input
 Attribute Source: vri

Sequence:	39
Optional:	Y
Format:	number
Length:	2
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: vertical_complexity Short Name: vert_compl

Alias vertical_complexity

Forestry Term: Vertical Complexity

Description: The subjective classification that describes the form of each tree layer as indicated by the relative uniformity of the height of the forest canopy as it appears on mid-scale aerial photographs. Vertical complexity is influenced by stand age, species (succession as it relates to shade tolerance) and degree and age of past disturbances. The tree height range is calculated as the total difference in height between the tallest and shortest visible dominant, co-dominant, and high intermediate trees. To most adequately represent the tree layer of interest, occasional occurrences of either very tall or very short trees should be ignored so that the vertical complexity indicated is for the majority of stems in the dominant, co-dominant, and high-intermediate portion of each tree layer. Vertical complexity is a subjective classification that describes the form of each tree layer as indicated by the relative uniformity of the forest canopy as it appears on mid-scale aerial photographs.

Measurement Criteria Vertical complexity is influenced by stand age, species (succession as it relates to shade tolerance) and degree and age of past disturbance.

Standard: Calculate the percent difference in tree height for the assignment of the Tree Vertical Complexity code.

Default:

Permitted Values Codes Description

1 Very uniform

A very uniform canopy with less than 11% difference between the height of the leading species and the average tree layer height. Holes (or canopy gaps) are generally not visible in the canopy and there is usually no evidence on the photograph of recent disturbances affecting the form of the stand. Examples include plantations and young, immature stands of shade intolerant species.

2 Uniform

A uniform canopy with 11% - 20% difference between the height of the leading species and the average tree layer height. A few holes (or canopy gaps) may be visible in the canopy and there is usually little or no evidence on the photograph of recent disturbance affecting the form of the stand.

3 Moderately uniform

A moderately uniform canopy with 21% - 30% difference between the height of the leading species and the average tree layer height. Some holes (or canopy gaps) may be visible in the canopy and there may be evidence of past disturbance affecting the form of the stand. Stocking may be somewhat patchy or irregular. Examples include older spruce-balsam stands.

4 Non-uniform

A relatively non-uniform canopy with 31% - 40% difference between the height of the leading species and the average tree layer height. Holes (or canopy gaps) are often visible in the canopy (due to past disturbance) and stocking is typically patchy or irregular.

5 Very non-uniform

A very non-uniform canopy with more than a 40% difference between the height of the leading species and the average tree layer height. Stocking is typically very patchy or irregular. Examples include disturbed dry belt Douglas-fir stands and decadent, coastal over-mature stands.

Input Format: #
 Input Example: 4
 Data Origin: input
 Attribute Source: vri

Sequence:	38
Optional:	Y
Format:	number
Length:	1
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes: Vertical complexity is used to identify and describe even-age and uneven-aged stands for further analysis in forest stand management and wildlife habitat assessment.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: vri_dead_stems_per_ha Short Name: dead_stems
 Alias vri dead stems per ha

Forestry Term: VRI Dead Stems per Hectare

Description: The number of standing dead trees visible to the photo interpreter in the dominant, codominant and high intermediate crown layer. Snag frequency is expressed as stem per hectare for each tree layer. The snag frequency provides a direct estimate of snags per hectare that can be used for wildlife and fire management. Note: Dominant trees have well-developed crowns that extend above the general level of the trees around them. Codominant trees have crowns forming the general level of trees around them. High intermediate trees have smaller crowns slightly below but extending into the general level of trees around them.

Measurement Criteria The following is a suggested approach to estimating snag frequency:
 -Select representative areas of the polygon.
 -Estimate the number of snags per hectare for each tree layer in the polygon.

Standard: Snag frequency is expressed as stems per hectare for each tree layer.

Default:

Permitted Values

Input Format: ###
 Input Example: 750
 Data Origin: input
 Attribute Source: both

Sequence:	35
Optional:	Y
Format:	number
Length:	4
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes: The snag frequency provides a direct estimate of snags per hectare that can be used for wildlife and fire management and provides information for danger tree assessment.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type: veg_comp_layer

Attribute Name: vri_live_stems_per_ha Short Name: live_stems
 Alias vri live stems per ha

Forestry Term: VRI Live Stems per Hectare

Description: The average number of living trees visible to the photo interpreter in the dominant, co-dominant and high intermediate crown positions in each tree layer in the polygon. It is expressed as stems per hectare. This attribute is also called stand density.

Measurement Criteria Estimate the density of trees in the polygon for each tree layer to the nearest stem per hectare when practical.

Standard: It is expressed as stems per hectare.

Default:

Permitted Values

Input Format: ####
 Input Example: 2252
 Data Origin: input
 Attribute Source: both

Sequence:	34
Optional:	Y
Format:	number
Length:	6
Decimal Places:	
Null:	Y

Use:

Linkage:

Relationship:

Sub Type Links: veg_comp_layer

Notes: Density provides a direct estimate of tree stems per hectare.

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual

Sub Type:

Attribute Name: whole_stem_biomass_per_ha Short Name: bioms_ws
Alias whole_stem_biomass_per_ha

Forestry Term: Whole Stem Biomass

Description: this is the total whole stem biomass per hectare of all species on a utilization of 4.0cm expressed as tonnes/ha

Measurement Criteria

Standard:

Default:

Permitted Values

Input Format: ###.##
Input Example: 235.6
Data Origin: derived
Attribute Source: both

Sequence:	
Optional:	
Format:	numeric
Length:	8
Decimal Places:	0
Null:	

Use:

Linkage:

Relationship:

Sub Type Links:

Notes:

Tips and Hints:

Reference: Ministry of Forests and Range, Forest Analysis and Inventory Branch, Vegetation Resources Inventory, Photo Interpretation Procedures Manual